


DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

HC70A, SAS70A, & PLSS059 Winter 2019 Genetic Engineering in Medicine, Agriculture, and Law

Professors Bob Goldberg, John Harada, &
Channapatna Prakash


Lecture 5
The Nuts & Bolts of Genetic Engineering:
From Mutations to Pedigrees to Drug
The Factor XIII Story

UCLA

TUSKEGEE
UNIVERSITY

UC DAVIS
UNIVERSITY OF CALIFORNIA


THEMES



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting




Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow


1. What is Hemophilia and How is it Inherited?
2. How Can a Disease Gene Be Found When It is Not Known Where the Gene is Expressed?
3. What Vectors Can Be Used For Cloning DNA?
4. What is the Advantage of Using a Virus Vector For Constructing Genome Libraries?
5. How To Make a Library of the Human Genome?
6. How Find a Gene With Only a Knowledge of the Protein Sequence?
7. How Use DNA Testing to Detect Factor VIII Disease Alleles?
8. How Isolate a Factor VIII cDNA Clone?
9. Genomic vs. cDNA Libraries
10. How Produce Factor VIII Protein For Use as a Drug?



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

How Do We Treat a Genetic Disease? From Gene To Drug

Due to Mutations in a Different Class of Blood Proteins

The Molecular Genetics of Hemophilia


(Potentially Lethal Disease)

Hemophiliacs bleed because a defective gene deprives them of a key blood-clotting protein. The protein has now been made artificially by isolating the normal gene and then inserting it into cultured cells

by Richard M. Lawn and Gordon A. Vehar

A Case Study of Cloning Genes and mRNAs


Reference: Scientific American, March 1, 1986



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria





DNA Fingerprinting





Cloning: Ethical Issues
and Future Consequences




Plants of Tomorrow



Founded in 1976 By Robert Swanson and Herb Boyer
First IPO in 1980 for \$88/share
Purchased by Hoffmann-La Roche in 2009 for \$47B



Hemophilia Has Been Known As An Inherited Disease For >2500 Years!



First Reference to Hemophilia is in the Old Testament

Genesis 17:10-14

'This is My covenant that you shall keep between Me and you and your descendants after you: every male among you shall be circumcised. You shall circumcise the flesh of the foreskin.....At the age of eight days every male among you shall be circumcised throughout your generations.....an uncircumcised male...that soul shall be cut off from its people, he has invalidated My covenant.'



The Talmud also makes reference to families in whom children have died as a result of circumcision (Babylonian Talmud, Chapter Yevamoth p64b) [6].

should a mother lose two children or should two sisters lose a child each after circumcision, subsequent children of the woman, the two sisters or of any other sisters of the same family should not be circumcised until they are older, or possibly not at all. This is thought to be the earliest reference to haemophilia; it was recognized in the Talmud that this condition was transmitted by the mother.



Abraham was circumcised at 93 and gave birth to Isaac at 99. His wife - Sarah - was 90!

A Cascade Of Events After Wounding Leads to A Fibrin Clot



LIFE 8e, Figure 49.10 (Part 2)

LIFE: THE SCIENCE OF BIOLOGY, Eighth Edition © 2007 Sinauer Associates, Inc. and W. H. Freeman & Co.

Clotting factors:

1. Released from platelets and injured tissue
2. Plasma proteins synthesized in liver and circulated in inactive form

Prothrombin
circulating
in plasma

Thrombin

Fibrinogen
circulating
in plasma


Fibrin

LIFE 8e, Figure 49.10 (Part 1)

LIFE: THE SCIENCE OF BIOLOGY, Eighth Edition © 2007 Sinauer Associates, Inc. and W. H. Freeman & Co.

Clotting Factors Such As Factor VIII
Play A Critical Role in This Process

Protein Factors in Blood Lead To Clotting



CLOTTING CASCADE begins when cell damage at a wound somehow activates the enzyme factor XII; it ends with the conversion of fibrinogen into fibrin by thrombin. At each step an inactive protein is converted into a protease, or protein-cutting enzyme (*color*), which activates the next protein. Some steps require cofactors such as factors VIII and V. The cascade includes positive- and negative-feedback loops (*colored arrows*). Thrombin activates factors VIII and V; it also deactivates them (by activating protein C), which helps to halt clotting. Some 85 percent of hemophiliacs lack factor VIII. The rest lack factor IX.

Eight Proteins/Genes Required:

1. Factor VII
2. Factor XI
3. Factor IX
4. Factor VIII
5. Factor X
6. Protein C
7. Prothrombin
8. Fibrinogen

What Happens If Any of These Proteins, or Genes, are Mutated?



No Blood Clot!

Hemophiliacs Have Mutations in Factor VIII, Factor IX, or Factor XI Genes

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 13.2

Some Important Genetic Disorders

Disorder	Symptom	Defect	Dominant/ Recessive	Frequency Among Human Births
Cystic fibrosis	Mucus clogs lungs, liver, and pancreas	Failure of chloride ion transport mechanism	Recessive	1/2500 (Caucasians)
Sickle cell anemia	Blood circulation is poor	Abnormal hemoglobin molecules	Recessive	1/600 (African Americans)
Tay-Sachs disease	Central nervous system deteriorates in infancy	Defective enzyme (hexosaminidase A)	Recessive	1/3500 (Ashkenazi Jews)
Phenylketonuria	Brain fails to develop in infancy	Defective enzyme (phenylalanine hydroxylase)	Recessive	1/12,000
Hemophilia	Blood fails to clot	Defective blood-clotting factor VIII	X-linked recessive	1/10,000 (Caucasian males)
Huntington disease	Brain tissue gradually deteriorates in middle age	Production of an inhibitor of brain cell metabolism	Dominant	1/24,000
Muscular dystrophy (Duchenne)	Muscles waste away	Degradation of myelin coating of nerves stimulating muscles	X-linked recessive	1/3700 (males)
Hypercholesterolemia	Excessive cholesterol levels in blood lead to heart disease	Abnormal form of cholesterol cell surface receptor	Dominant	1/500


18,000 People in US Have Hemophilia & 400 Babies/Year Are Born With Disorder Prior to 1960s - Average Life Span Was 11 Years

Hemophilia A	Defective Factor VIII Gene	1/10,000 males	80%
Hemophilia B	Defective Factor IX Gene	1/30,000 males	20%
Hemophilia C	Defective Factor XI Gene	Autosomal	<1%

**Both Factor VIII & IX Genes
on X-Chromosome (♀ → ♂'s)**


Factor VIII and Factor IX Genes are Closely Linked on the X Chromosome

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Nature, March, 2005


The X chromosome has ~1098 Genes and 150,000,000 bp (150 Mb). **168 Mendelian Diseases Explained by 113 X-Linked Genes**



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

**Pedigrees Can Be Used To Determine If
a Trait is Dominant or Recessive**


**Each Type of Inheritance Predicts
Specific Results in Each Generation**

Hemophilia A and B Genes Are Sex Linked & Recessive Traits When Mutated

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© Bettmann/Corbis



- Note:**
1. Males Obtain Defective Gene From Mothers
 2. 50% of Sons Of A Maternal Carrier Have The Defective Gene

Hemophilia A and B Sex-Linked Inheritance

		Carrier Female	
		Egg	X
Healthy Male	Sperm		
	X	XX ♀ <i>Carrier</i>	XX ♀ <i>Healthy</i>
	Y	XY ♂ <i>Hemophiliac</i>	XY ♂ <i>Healthy</i>

Sex-Linked Inheritance

♀ Carriers → 1/2 Sons Afflicted + No Daughters!
 Only One X-Chromosome is in ♂

What Was Known About Factor VIII *Before Gene Cloned?*

1. Blood Protein (But Perhaps Synthesized Elsewhere!)
2. Could be purified in small amounts from >20 Liters of human blood + cow blood + pig blood
3. Short Stretch of Protein Sequenced = Known Protein Sequence!
4. Hemophilia A could be treated by blood transfusions from normal individuals, ∴ clotting factor in blood
5. 1980s AIDS Epidemic Caused Many Hemophiliacs to Get HIV/AIDS (~50% of hemophiliacs got AIDS in 1985)


∴ How to go From Protein to Gene

The Problem

For Factor VIII- Not Known Where Gene Was Expressed ∴ **Must Use Genome Library**

Early 1980's


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Key Concept



How Clone A Gene When You Don't Know Where it is Expressed !



Key:
Protein Sequence Known



How Find Gene & cDNA?

Protein → Gene → mRNA → Drug !

mRNA → Drug


© 2001 Terese Winslow (assisted by Lydia Kibiuk)

Steps Required to Clone Factor VIII Gene and cDNA

Gene

cDNA

1. Make Genome Library Because Factor VIII Gene in Genome!
2. Purify Protein from Blood- that's where it works (wasn't known where made)
3. Reverse Translate using the genetic code a portion of the protein sequence
4. Synthesize a DNA probe complementary to Factor VIII gene corresponding to protein sequence
5. Screen Genome Library Entire Gene on The Clone?




1. Use Gene probe to screen cDNA library for Factor VIII cDNA clone
 2. How know what mRNA to use to make cDNA library?
 3. Use gene probe to probe RNA blots from all major organs (liver, kidney, blood, etc.)
 4. Find Factor VIII mRNA in liver-male, liver- secrete into blood
- Why Need cDNA?**
Story continued

Want cDNA to Manufacture Factor VIII as a Drug to Treat Hemophilia A!

Step One

How to Construct a Human Genome Library to Find the Factor VIII Gene?


If It is Not Known Where Gene is Active
Can "Look" to Genome Instead of mRNA to
Find + Clone Gene!



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

Vectors Used in Genetic Engineering Have Similar Conceptual Properties But are Used in Different Situations

Table 3.2 A COMPARISON OF DNA VECTORS AND THEIR APPLICATIONS

Vector Type	Maximum Insert Size (kb)	Applications	Limitations
Bacterial plasmid vectors (circular)	~6-12	DNA cloning, protein expression, subcloning, direct sequencing of insert	Restricted insert size; limited expression of proteins; copy number problems; replication restricted to bacteria
Bacteriophage vectors (linear)	~25	cDNA, genomic and expression libraries	Packaging limits DNA insert size; host replication problems
Cosmid (circular)	~35	cDNA and genomic libraries, cloning large DNA fragments	Phage packaging restrictions; not ideal for protein expression; cannot be replicated in mammalian cells
Bacterial artificial chromosome (BAC, circular)	~300	Genomic libraries, cloning large DNA fragments	Replication restricted to bacteria; cannot be used for protein expression
Yeast artificial chromosome (YAC, circular)	200-2,000	Genomic libraries, cloning large DNA fragments	Must be grown in yeast; cannot be used in bacteria
Ti vector (circular)	Varies depending on type of Ti vector used	Gene transfer in plants	Limited to use in plant cells only; number of restriction sites randomly distributed; large size of vector not easily manipulated


Copyright © 2009 Pearson Education, Inc.

Properties of All Vectors

1. Replicate
2. Selectable
3. Can Be Used To Insert Foreign Genes/Restriction Sites
4. Easily Isolated + Transferred Back To Cells


Plasmid vs. Bacteriophage Vectors for Cloning DNA Fragments

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



a.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




b.




**“Natural”
Infection Process**

- **Much More Efficient**
- **Can Use Less DNA**
- **Get Lots More Clones**
- **Need Lots of Clones For Large Genome**

Structure of the λ Phage and Its Genome




(b) λ Phage assembly



First Genome Sequence


Sanger et al. 1982 *J. Mol. Biol.*
162: 729-773.



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting





Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

λ Phage Infects *E.coli* & Destroys (Lyses) Cells






DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting




Cloning: Ethical Issues
and Future Consequences




Plants of Tomorrow

Lysed Cells Can Be Seen as Clear Plaques on Agar Plates



(a)




(b)

1. Each Plaque is a Virus
Clone Representing One
Viral Infection!

2. Selectable Marker is
Bacterial Cell Destruction &
Plaque Formation

Advantages of λ Virus as a Vector for Cloning DNA


1. Long DNA Segments Can Be Cloned (~20kb) Need Fewer Clones For Whole Genome!
2. Can Clone DNA Segments In Viral Genome & Self-Assemble With Viral Proteins Into Virus In A Test Tube!
∴ Make Recombinant Viruses In The Lab!
3. Use “Natural” Infection Process To Generate Large Number Of Clones For A Eukaryotic Genome Library.
Much Higher Efficiency For Getting Recombinant DNA
→Bacterial Cells Compared With Dna Transformation.
∴ Set More Clones Per Amount Of Recombinant DNA!



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting




Cloning: Ethical Issues
and Future Consequences




Plants of Tomorrow

Using a Bacterial Virus To Clone the Human Genome



(b)



Copyright © 2009 Pearson Education, Inc.

Mixture of Plaques =
Library With All Human
DNA Sequences
Represented

Cloning the Human Genome and Screening for the Factor VIII Gene







Figure 20-6
 Introduction to Genetic Analysis, Ninth Edition
 © 2008 W. H. Freeman and Company

Why Partial Digestion? An Important Concept!
What is Complete & Partial Digestion?

Partial Digestion Permits "Walking" From One DNA Region to the Next



Constructing a Human Genome Library by Partial Digestion Creates a Set of Overlapping DNA Fragments/ Clones



Step Two

**How Find the Factor VIII
Gene in a Human
Genome Library?**

The Genetic Code


		Second Letter					
		U	C	A	G		
1st letter	U	UUU Phe UUC UUA Leu UUG	UCU UCC Ser UCA UCG	UAU Tyr UAC UAA Stop UAG Stop	UGU Cys UGC UGA Stop UGG Trp	U C A G	
	C	CUU CUC Leu CUA CUG	CCU CCC Pro CCA CCG	CAU His CAC CAA Gln CAG	CGU CGC Arg CGA CGG	U C A G	
	A	AUU AUC Ile AUA AUG Start Met	ACU ACC Thr ACA ACG	AAU Asn AAC AAA Lys AAG	AGU Ser AGC AGA Arg AGG	U C A G	
	G	GUU GUC Val GUA GUG	GCU GCC Ala GCA GCG	GAU Asp GAC GAA Glu GAG	GGU GGC Gly GGA GGG	U C A G	

Properties

- Universal
- Three Nucleotides
- Punctuation
- Degenerate

Factor VIII Protein → Gene

Using the Factor VIII Protein Sequence and Genetic Code as a Guide to Synthesize a Factor VIII Probe



2. Make Several Probes All Codon Combinations!

3. One Will Be Correct Probe

1. Use Genetic Code


How many Combinations of Synthetic Probes?

$$2 \times 3 \times 2 \times 1 \times 2 \times 2 \times 2 = 96$$

Using the Genetic Code to go From Protein Sequence to Gene Sequence

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


(b) Synthesizing DNA probes based on reverse translation




1. Need Amino Acid Sequence of Part of the Protein
2. Need DNA Sequences Representing all Codon Combinations
3. Synthesize DNA Sequence Probes!

Finding The Factor VIII Gene Or Part of Gene!!


Human Genome Library Clones





Transfer plaque to absorbent membrane.



Incubate membrane with radioactive probe.



Autoradiograph to locate desired clone.



Master Plate

Infect fresh bacterial host.



Purify Factor VIII Genomic Clone

Amplify desired gene.

Sequence To See If it Matches Probe/Protein






Fig. 1 Detection of the factor VIII gene with probe 8.3, 5'-CTTTCCAGGTCAACGTCGGAGAAATAAGCCCAAGC, a



: Analysis, Ninth Edition
rd Company

The Result-The Factor VIII Gene is Huge- 186,000 bp- The Probe Identified a Clone Containing **Only One Part of Gene !!!** Why?



How Find Clones with Rest of Gene?

Key Question !


Remember - the Library Contains Overlapping DNA Clones \therefore Can Use One Part of First Clone to Re-Screen Library & "Walk" to Other Gene Regions - Using Restriction Maps & Sequencing (Compare With Protein Sequence) as Guides!

Sequence -----> GenBank

Step Three

Finding the Entire Factor VIII Gene? Walking & Sequencing

Walking Up and Down Genes and Chromosomes



**Reiterative Library
Screening Process**

**Find Overlapping
Clones By
Restriction Site
Mapping**

Figure 20-13
Introduction to Genetic Analysis, Ninth Edition
© 2008 W. H. Freeman and Company

Basis of Genome Projects & Whole Genome Sequencing


Key
Concepts

How know Find Complete Factor VIII Gene?

Compare Protein & DNA Sequences

The Factor VIII Gene Was Found To Be Very Large

- **186,000 Nucleotides in Length** (Won't Fit in One Phage Clone)
- **25 Introns**
- **9,000 Nucleotide Coding Sequence (cDNA)**
- **2,351 Amino Acids in Protein**



Factor VIII Mutations Occur Throughout the Gene

[*Haemophilia* 11, 481-491 (2005)]


Factor VIII gene mutations in haemophilia A patients without intron 22 inversion.

VIII:C (%)	Family history	Consanguinity*	Inversion	Codon†	Mutation	Amino acid change	Exon	Conservation‡
1	Sporadic	NC	Normal	51	TTT → TCT§	Phe → Ser	2	FFFF, identical
1.20	Sporadic	NC	Normal	80	GTT → GAT	Val → Asp	3	VVVV, identical
1	Sporadic	NC	Normal	102	GGT → GTT§	Gly → Val	3	GGGG, identical
2	Sporadic	NC	Normal	104	TCC → CCC§	Ser → Pro	3	SSSS, identical
6	Sporadic	NC	Normal	143	GAG → AAG§	Glu → Lys	4	EEEE, identical
1	Sporadic	NC	Normal	233	delCA§	Thr → fs (TGA-264)	6	
2.70	Inherited	NC	Normal	321	GAA → AAA	Glu → Lys	8	EEEE, identical
0	Sporadic	NC	Normal	372	CGC → CAC	Arg → His	8	RRRR, identical
3	Inherited	NC	Normal	527	CGG → TGG	Arg → Trp	11	RRRR, identical
1	Sporadic	NC	Normal	528	TGC → TAC§	Cys → Tyr	11	CCCC, identical
1	Inherited	NC	Normal	592	CAA → TAA	Gln → Stop	12	QQQQ, identical
1	Inherited	NC	Normal	864	delGACA insCAATTAAATGAGAA§	Gly → fs [TAA-867]	14	
1	Sporadic	NC	Normal	948	insA§	Lys → fs (TGA-984)	14	
1	Sporadic	NC	Intron 1	1107	AGG → TGG§	Arg → Trp	14	RGKK, dissimilar
1	Sporadic	NC	Normal	1107	AGG → TGG§	Arg → Trp	14	RGKK, dissimilar
1	Inherited	NC	Normal	1191-1194	delA	Ile → fs (TAG-1198)	14	
1.40	Sporadic	NC	Normal	1191-1194	insA	Ile → fs (TAA-1220)	14	
1	Sporadic	C	Normal	1227	delC§	Leu → fs (TGA-1231)	14	
2.10	Sporadic	NC	Normal	1241	GAC → GAG	Asp → Glu	14	DGGE, similar
1	Sporadic	NC	Normal	1392	1392del1418§	Pro → fs (TAG-1446)	14	
1	Inherited	C	Normal	1392	1392del1418§	Pro → fs (TAG-1446)	14	
1	Sporadic	NC	Normal	1441	insA§		14	
1	Inherited	C	Normal	1441	insA§		14	
1	Inherited	NC	Normal	1502	CAG → TAG§	Gln → Stop	14	QREQ, dissimilar
1	Inherited	NC	Normal	1504	delGT§	Val → fs (TGA-1517)	14	
1	Sporadic	NC	Normal	1535	TGG → TGA	Trp → Stop	14	WLWM, dissimilar
inhibitor 96 BU								
1	Sporadic	NC	Normal	1571	TAT → TAA§	Tyr → Stop	14	Y-YY, dissimilar
1	Sporadic	NC	Normal	1581	AAA → TAA§	Lys → Stop	14	KEKK, dissimilar
0.20	Sporadic	NC	Normal	1696	CGA → GGA	Arg → Gly	14	RRRR, identical
1.80	Sporadic	NC	Normal	1729	delA§	Gln → fs (TAA-1752)	15	
1	Inherited	NC	Normal	1751	GAA → AAA§	Glu → Lys	15	EEEE, identical
1	Sporadic	NC	Normal	1775	TTC → TCC§	Phe → Pro	16	FFFF, identical
1	Sporadic	NC	Normal	1835	TGG → TGA§	Trp → Stop	16	WWWW, identical
7.60	Sporadic	C	Normal	1882	ATC → ATA§	Ile → Ile	17	III, identical
3	Inherited	C	Normal	1966	CGA → CAA	Arg → Glu	18	RRRR, identical
1	Sporadic	NC	Normal	1966	CGA → TGA	Arg → Stop	18	RRRR, identical

FVIII GENE MUTATIONS IN INDIAN PATIENTS

Need To Screen Across the Gene for Markers -- Family Specific

Factor VIII Gene Probes/ Sequence Can Be Used to Characterize Mutant Genes & Do DNA Testing for Carriers



Mutations in Factor Gene



Once Gene & cDNA Identified!


Use DNA Gel Blots (or PCR) & Factor VIII Probes to Investigate Presence of Mutant Alleles in Families (carriers)

Mutations Arise Independently in Families

Using PCR and RFLPs (Markers) to Screen For the Hemophilia A Disease Allele/Gene

1. Use PCR To Amplify A Specific Factor VIII Gene Region
2. Use Restriction Enzyme (Bcl I) To Distinguish Between Normal Allele (1 Site) & Disease Allele (No Site)


= = Normal Allele
- = Disease Allele



The 21st Century Approach!

1. Sequence the Entire Gene & Find Mutation
2. Then Synthesize Primers to Test Family Members Using PCR


Only Can Do This With a Knowledge of DNA Sequence of Wild-type (Normal) and Disease Genes (Can Vary Family to Family)



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

Step Four

How Find Factor VIII mRNA to
Generate a cDNA for Protein
Production in Host Cells?


Recall: Eukaryotic Genes Provide
Obstacles for Efficient Protein
Production in Genetically
Engineered Cells!

Introns! Switches!


Making the Drug

Need cDNA Not Gene

Factor VIII Gene Can Be Used to Find Out Where It is Active Using RNA Blots




4. Wash away unhybridized probe. Make autoradiograph.



RNA Blot Is Like a DNA Blot Except That RNA is on Gel & Blotted


3. Separate RNA samples by gel electrophoresis. Blot onto filter. Expose filter to labeled hybridization probe.




Factor VIII Gene Is Highly Active in Liver!

Could Also Use PCR (RT-PCR)

Using Factor VIII Gene Probe to Identify Factor VIII cDNA clone






DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting




Cloning: Ethical Issues
and Future Consequences




Plants of Tomorrow

Genomic Libraries vs. cDNA Libraries A Review




Engineer Factor VIII cDNA to Produce Protein in Host Cell & Synthesize Factor VIII in Mammalian Cells


(a) Transient transfection



(b) Stable transfection (transformation)




Why Mammalian Cells?



Purify Factor VIII Protein!

Need FDA Approval Before Recombinant DNA Drug Can Be Marketed and Used to Treat Patients



A Long and Expensive Process!

Recombinant Factor VIII



Bayer Biological Products EU



Bayer HealthCare

Biological Products Division

Search | Sitemap

Home About Us **About Haemophilia** For Kids Research & Development Press Releases

→ **Recombinant Factor VIII**

More Resources

Haemophilia Centres in Europe

Related Links

Haemo-QoL Project
Hemophilia Research Awards

Recombinant factor VIII

Recombinant factor VIII (rFVIII) is the antihemophilic factor A, obtained using recombinant DNA technology. With this technology, pure protein is synthesized in the laboratory instead of being extracted from blood. In the following pages, it will be explained in detail how the knowledge and analysis of DNA, using the new instruments of molecular genetics, have represented both the beginning and follow-up stages in the development of recombinant FVIII.




Prophylactic Treatment Costs \$300,000/Year! Most Hemophiliacs Use "On Demand" or As Needed

Factor VIII Gene Cloned In 1983
Factor VIII (Recombinant) Approved As Drug In 1993! Ten Years From Gene → Drug! (Off Patent In 2011)

Using Factor VIII to Treat Hemophilia

Formation of a Blood Clot



FIBRIN STRANDS stabilize a blood clot at the site of a wound by trapping the platelets that form the bulk of the clot. The electron micrograph, which was made by Jon C. Lewis of Wake Forest University, shows a clot formed in a suspension of platelets and fibrin.

A clot in the bloodstream is the result of a complex cascade of enzymatic reactions culminating in the conversion of fibrinogen, a soluble protein, into insoluble fibrin strands. In hemophiliacs a crucial protein in the blood-clotting cascade is either missing or defective.

A Triumph of Genetic Engineering

The Future: Gene Therapy - A Permanent "Cure"

December 10, 2011

Treatment for Blood Disease Is Gene Therapy Landmark

By NICHOLAS WADE


TIME Partners
with
ON.

Gene Therapy Shows Promise for Treating Hemophilia

By ALICE PARK Monday, December 12, 2011

The First Ever In-Human Gene Editing Will Try and Combat Hemophilia


Factor IX - Hemoglobin B
FDA-Approved Clinical Trial
2016



DNA
Genetic Code of Life



Entire Genetic Code
of a Bacteria



DNA Fingerprinting



Cloning: Ethical Issues
and Future Consequences



Plants of Tomorrow

The Factor VIII Story -- A Summary

1. Purify Small Amounts of Factor VIII
2. Obtain Partial or Complete Amino Acid Sequence
3. Use the Genetic Code to Synthesize Degenerate DNA Probes
4. Isolate Factor VIII DNA Clones Complementary to Probe in Genome Library
5. Determine if Factor VIII Clones Contain the Complete Gene By Sequencing and Comparing With Protein Sequence
6. If Not, "Walk" to Obtain Overlapping DNA Clones That Collectively Contain the Factor VIII Gene
7. Sequence Clones To Determine Where the Factor VIII Gene Starts and Stops
8. Use Factor VIII Genome Probe to Find Out What Body Organ/Tissue Expresses the Factor VIII Gene
9. Make a cDNA Library From the Target Organ/Tissue and Isolate a Factor VIII cDNA Clone
10. Sequence the Factor VIII cDNA Clone and Compare With Factor VIII Gene Sequence to Map its Anatomy (I.e., introns, exons, swtiches) and Ensure That it Contains the Complete Protein Coding Sequence
11. Use Factor VIII cDNA and/or Genome Fragments as a Probe to Find RFLP Markers For Disease Alleles -- Or Sequence Disease Alleles to Find Relevant RFLP Markers By Comparison With Wild-Type Sequence
12. Insert Factor VIII cDNA Into an Expression Vector and Synthesize Factor VIII Protein in Host Cells (e.g., Mammalian Cells)