

Developmental Cell

Multi-level Modulation of Light Signaling by GIGANTEA Regulates Both the Output and Pace of the Circadian Clock

Highlights

- The clock component GI gates light signaling through direct regulation of PIFs
- GI modulates PIF transcriptional activity via several distinct molecular mechanisms
- Gating of PIF activity by GI is required to time key output processes such as growth rate
- Regulation of PIF activity by GI is required for optimal circadian clock progression

Authors

Maria A. Nohales, Wanlu Liu, Tomas Duffy, ..., Julin N. Maloof, Steven E. Jacobsen, Steve A. Kay

Correspondence

stevekay@usc.edu

In Brief

Nohales et al. show how the clock component GIGANTEA modulates light signaling pathways by gating the activity of PIFs at multiple regulatory levels. This function of GI is essential not only to optimize growth strategies in plants but also to set the pace of the clock in response to light-dark cycles.



Multi-level Modulation of Light Signaling by GIGANTEA Regulates Both the Output and Pace of the Circadian Clock

Maria A. Nohales,¹ Wanlu Liu,^{2,3} Tomas Duffy,¹ Kazunari Nozue,⁴ Mariko Sawa,⁵ Jose L. Pruneda-Paz,⁵ Julin N. Maloof,⁴ Steven E. Jacobsen,^{2,6} and Steve A. Kay^{1,7,*}

¹Keck School of Medicine, University of Southern California, Los Angeles, CA 90089, USA

²Department of Molecular, Cell, and Developmental Biology, University of California, Los Angeles, Los Angeles, CA 90095, USA

³Zhejiang University, University of Edinburgh Institute, Zhejiang University School of Medicine, Hangzhou 310058, P.R. China

⁴Department of Plant Biology, University of California, Davis, Davis, CA 95616, USA

⁵Section of Cell and Developmental Biology, Division of Biological Sciences, University of California, San Diego, San Diego, La Jolla, CA 92093, USA

⁶Howard Hughes Medical Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

⁷Lead Contact

*Correspondence: stevekay@usc.edu

<https://doi.org/10.1016/j.devcel.2019.04.030>

SUMMARY

Integration of environmental signals with endogenous biological processes is essential for organisms to thrive in their natural environment. Being entrained by periodic environmental changes, the circadian clock incorporates external information to coordinate physiological processes, phasing them to the optimal time of the day and year. Here, we present a pivotal role for the clock component GIGANTEA (GI) as a genome-wide regulator of transcriptional networks mediating growth and adaptive processes in plants. We provide mechanistic details on how GI integrates endogenous timing with light signaling pathways through the global modulation of PHYTOCHROME-INTERACTING FACTORS (PIFs). Gating of the activity of these transcriptional regulators by GI directly affects a wide array of output rhythms, including photoperiodic growth. Furthermore, we uncover a role for PIFs in mediating light input to the circadian oscillator and show how their regulation by GI is required to set the pace of the clock in response to light-dark cycles.

INTRODUCTION

Accurate decoding of environmental signals and integration of these cues within cellular networks are essential for organisms to succeed in their natural environment. The rhythmic and periodic nature of relevant external conditions, such as light and temperature oscillations, has driven the evolution of endogenous molecular oscillators that enable organisms to anticipate these cyclic changes and coordinate key physiological processes accordingly (Millar, 2016). In plants, adequate integration of environmental cues and precise phasing of biological processes are

key, as their sessile growth habit precludes their escape from disadvantageous conditions. The influence of the circadian clock on plant development is pervasive. Multiple processes, including growth, stress responses, and developmental transitions, are coordinated by the clock in conjunction with other signaling pathways (Greenham and McClung, 2015; Sanchez and Kay, 2016). Clock genes and their feedback regulatory mechanisms have been extensively studied in plants (Nohales and Kay, 2016). However, little is known about how environmental information is transmitted to this complex network nor do we have mechanistic knowledge on how the clock regulates such a wide array of biological processes.

A key clock protein that seems to function at the interface between the oscillator and its output is GIGANTEA (GI), a conserved plant-specific protein expressed in the evening (Fowler et al., 1999; Park et al., 1999). GI is essential for accurate timekeeping and clock synchronization with the environment (Gould et al., 2006; Kim et al., 2007; Locke et al., 2006; Martin-Tryon et al., 2007; Mizoguchi et al., 2005). Besides its role in the central oscillator, GI modulates myriad clock output pathways, including abiotic stress (Cao et al., 2005; Kim et al., 2013a), photoperiodic flowering (Sawa et al., 2007; Suárez-López et al., 2001), and light signaling (Huq et al., 2000; Martin-Tryon et al., 2007; Oliverio et al., 2007). Despite its pivotal role in clock function and plant development, knowledge of the mechanisms by which GI is able to influence such a wide array of cellular networks is only starting to emerge. At the post-translational level, GI interacts with multiple proteins from diverse pathways (Mishra and Panigrahi, 2015) and, recently, a role for GI as a co-chaperone (holdase) has been uncovered (Cha et al., 2017). In the context of transcriptional regulation, GI has been shown to influence the transcription of flowering time genes through interaction with and modulation of their transcriptional regulators, as well as by occupancy of a small cluster of promoter regions (Sawa and Kay, 2011; Sawa et al., 2007).

Light signaling entails the perception of light quality and quantity by an array of photoreceptors specialized in sensing specific wavelengths of the light spectrum (Möglich et al.,



2010), which then relay this information to transcriptional networks to ultimately regulate the expression of genes involved in light responses. In terms of light quality, the red and far-red regions are especially relevant and are perceived by the phytochrome (phy) family of photoreceptors (Xu et al., 2015). One of the mechanisms through which phytochromes achieve regulation of gene expression is through interaction with PIFs, a family of basic helix-loop-helix (bHLH) transcription factors that function as negative regulators of photomorphogenesis in the dark (Leivar and Monte, 2014; Xu et al., 2015). Notably, these factors carry out a broader function and act as hubs that integrate information from multiple cellular pathways, including light, temperature, hormone, and circadian signaling (Castillon et al., 2007; Legris et al., 2017; Leivar and Quail, 2011).

Previous studies have uncovered extensive connections between the oscillator and PIFs, and several clock components have been identified to regulate *PIF* expression and/or activity (Martín et al., 2018; Nieto et al., 2015; Soy et al., 2016; Zhu et al., 2016). Here, we have investigated GI function in the context of transcriptional regulation and have uncovered direct connections of unprecedented complexity between this core clock component and PIFs. We show how GI globally modulates light signaling by gating the activity of the PIF proteins at multiple regulatory levels and, at the physiological level, we provide evidence regarding how this regulation influences output rhythms such as photoperiodic growth. Since PIF proteins function as hubs in the regulation of plant growth and development (Davière and Achard, 2016; Leivar and Quail, 2011; Xu et al., 2014), broad modulation of their activity by GI provides a mechanism by which the circadian clock acts as an overarching hub for integrating environmental and endogenous signaling pathways to optimize adaptive traits in plants. Moreover, we uncover a critical role for the PIF proteins in light input to the circadian system and show that their regulation by GI is required for optimal clock progression, providing a molecular framework for understanding how light signaling entrains the plant circadian clock.

RESULTS

GI Shares Targets with Both Clock and Light Signaling Pathways

We initiated our study of the molecular mechanisms of GI function in the context of transcriptional regulation by performing a reanalysis of published genome-wide gene expression datasets. A search for over-represented *cis* elements at the promoter regions of differentially expressed genes (DEGs) in the *gi-2* null mutant (Kim et al., 2012) revealed the G-box (CACGTG) and the evening element (EE) ([A]AAATATCT) as the two most highly enriched motifs (Figure 1A). The G-box motif is found in the promoters of many light regulated genes and is known to be bound by different transcription factors including the PIFs (Hornitschek et al., 2009; Huq and Quail, 2002; Martínez-García et al., 2000), whereas the EE is bound by the morning core clock repressors CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY) (Alabadi et al., 2001) and is enriched at the promoters of genes with peak expression in the evening (Covington et al., 2008; Harmer et al., 2000).

Given the involvement of GI in light signaling and photomorphogenesis (Huq et al., 2000; Martin-Tryon et al., 2007) as well

as in *CCA1* and *LHY* expression (Fowler et al., 1999; Mizoguchi et al., 2002; Park et al., 1999), we decided to subsequently analyze the overlap between DEGs in *gi-2* (Kim et al., 2012) and a comprehensive set of PIF-regulated genes (Oh et al., 2012) or DEGs in *cca1-1;lhy-11* (Kamioka et al., 2016). In spite of the diversity of conditions under which these legacy datasets were obtained, this analysis rendered two subsets of significantly shared genes with distinct characteristics (Figures 1B and S1A). The subset of genes shared by GI and PIFs is most highly enriched in Gene Ontology (GO) terms related to photosynthesis, light signaling, and response to abiotic stress and is enriched in peak expression phases around dawn (Zeitgeber time 23 (ZT23) to ZT1) and morning/early afternoon (ZT2 to ZT5), especially under short-day (SD) conditions (Figures 1C, S1B, and S1D). In contrast, genes shared by GI and *CCA1* and *LHY* are involved in pathways related to circadian rhythms, temperature acclimation, response to diverse abiotic stimuli, and response to gibberellins (Figure S1D). In this case, analysis of time-of-day expression further revealed that these genes tend to be expressed in the evening, mostly at ZT11 to ZT14 (Figure S1C).

These results support the notion that GI plays dual roles in the regulation of light signaling and the circadian system, consistent with earlier genetic analyses (Martin-Tryon et al., 2007; Mizoguchi et al., 2005; Oliverio et al., 2007). Genes shared by GI and *CCA1/LHY* are clock components and clock outputs whose evening-phased expression is likely driven by the EE, supporting a key role for GI in the central oscillator and in the regulation of *CCA1* and *LHY* amplitude (Fowler et al., 1999; Mizoguchi et al., 2002; Park et al., 1999). Conversely, GI and PIFs seem to intersect at a subset of genes for the regulation of photosynthesis, light signaling, and growth promotion at the end of the night.

GI Interacts with PIFs

GI acts as a positive regulator of light signal transduction and photomorphogenesis (Huq et al., 2000; Martin-Tryon et al., 2007), and *gi* mutant plants display a long hypocotyl phenotype under different light conditions (Figure S1E). While GI affects the expression of several clock factors, which in turn interact with and regulate the function of PIFs (Martín et al., 2018; Nieto et al., 2015; Soy et al., 2016; Zhu et al., 2016), we wondered if GI function in the light signaling pathway could also arise from a direct connection to PIFs. Given the ability of GI to interact and modulate the activity of diverse proteins, we hypothesized that interaction with PIF proteins may provide a molecular framework for understanding the mechanism by which GI directly regulates light signaling transcriptional networks. Leveraging an arrayed *Arabidopsis* transcription factor library (Pruneda-Paz et al., 2014), we performed a high-throughput yeast two-hybrid screen using GI as bait. We found that GI is able to strongly interact with PIF3, and to a lesser extent with other PIFs such as PIF1, PIF4, and PIF5, in this system (Figures S1F and S1G).

Through *in vitro* pull-down assays using full-length and deleted protein fragments, we verified the observed interactions (Figure 1D) and mapped the interaction domains for PIF3 (Figure S1H). These deletion studies showed that GI likely interacts with the bHLH DNA-binding motif of PIF3, suggesting that the interaction could hinder PIF3 binding to chromatin. We further confirmed the interaction of GI with PIF3 and PIF5 *in vivo* by




Figure 1. GI Shares Targets with Light Signaling Components and Interacts with PIF Proteins

(A) Over-represented *cis* motifs at the promoter regions of *gi-2* DEGs.
(B) Overlap between DEGs in *gi-2* and a comprehensive set of PIF-regulated genes (intersection $p < 2.2 \times 10^{-16}$).
(C) Phase enrichment heatmap depicting the p value of the phase of peak expression enrichment (count/expected) of genes differentially expressed in *gi-2* only (GI), regulated by PIFs only (PIFs), and potentially regulated by GI and PIFs (GI-PIFs) under SD conditions (* $p < 0.01$). Day period is marked in yellow and night period in gray.
(D) *In vitro* pull-down assays showing the interaction between GI and PIFs (PIF1, PIF3, PIF4, and PIF5).
(E) *In vivo* co-immunoprecipitations in *Arabidopsis* transgenic seedlings expressing GI-YPET-FLAG and PIF3-ECFP-HA (left panel) and GI-YPET-FLAG and PIF5-HA (right panel) tagged protein versions.
See also Figure S1 and Table S1.

performing co-immunoprecipitation (coIP) studies in transgenic *Arabidopsis thaliana* seedlings expressing tagged protein versions (Figure 1E).

GI Modulates PIF Stability and Activity

Given the physical interaction between GI and PIFs through a region containing the bHLH domain, we hypothesized that GI could function to prevent the access of PIFs to their target promoters and/or affect their stability. In fact, gene-expression analyses of PIF target genes revealed that these are significantly altered in *gi-2* and *Glox* plants across the entire day and even more significantly during the night period (Figures 2A and S2A). Although GI affects *PIF4* and *PIF5* mRNA expression during the early night (de Montaigu et al., 2015; Fornara et al., 2015) (Figure S2B), this observation seems insufficient to explain the dramatic effect observed on PIF target gene expression.

Transient stability analyses revealed a modest effect of GI on HA-PIF3 protein accumulation (Figure S2C), which was not evident for HA-PIF5 (Figure S2D). More significantly, we observed that GI co-expression promoted a shift in the migration pattern of both PIFs suggestive of post-translational modification. A treatment with phosphatase was observed to reverse the mobility shift (Figure S2E), indicating that it is dependent

upon phosphorylation. Because PIF phosphorylation is linked to their degradation (Ni et al., 2013; Shen et al., 2008), the results globally suggested that GI may affect PIF accumulation *in vivo*. Subsequent analysis of PIF5 protein accumulation at ZT8 and 16 in the *Arabidopsis* lines expressing tagged versions of GI and PIF5 confirmed that PIF5-HA accumulation is strongly affected by GI overexpression *in vivo* (Figures 2B and S2F).

We next performed transient transcriptional activation assays and measured activation of the *pPIL1::LUC* reporter (promoter of the well-characterized PIF target gene *PHYTOCHROME INTERACTING FACTOR 3-LIKE 1* (*PIL1*) driving the expression of the luciferase gene) as a proxy for PIF activity in the presence and absence of GI. In these experiments, PIF overexpression resulted in activation of the reporter as expected (Figures 2C and S2G), except in the case of PIF3 for which activation could not be assessed under our experimental conditions. In all cases, co-expression of GI together with the PIF effectors led to a drastic reduction in *pPIL1* promoter activation (Figures 2C and S2G). Given that GI affects PIF protein levels, we next examined whether the observed effect of GI on PIF transcriptional activity may also arise from a sequestration mechanism, similar to the DELLA proteins (de Lucas et al., 2008; Feng et al., 2008; Li et al., 2016). To test this hypothesis, we performed *in vitro*




Figure 2. GI Modulates PIF Stability and Activity

(A) Relative expression of *PIL1* and *AT5G02580* in wild-type (Col-0), *gi-2*, and Glox seedlings grown for 10 days in SDs (mean $\pm$ SEM of 3 biological replicates). White and gray shadings represent day and night, respectively.

(B) PIF5-HA protein accumulation at ZT8 and ZT16 in the indicated backgrounds under SD conditions. Protein levels were normalized against ACTIN levels. The quantitation is shown on the right (mean $\pm$ SEM of 3 biological replicates).

(C) Transactivation assays in *N. benthamiana* leaves. Different effectors were coexpressed with the *pPIL1::LUC* reporter construct. Luminescence was measured 2 days post infiltration in SD conditions. Results show mean $\pm$ SEM (n = 12).

(D and F) PIF3 (D left panel and F) and PIF5 (D right panel) ChIP assays of 10-day-old seedlings grown in SD conditions and harvested at the indicated ZTs. The enrichment of the specified regions in the immunoprecipitated samples was quantified by qPCR. Values represent mean $\pm$ SEM (n = 2-4).

(E, G, and H) Relative expression of *PIL1* in 10-day-old seedlings of the indicated backgrounds in SDs (mean $\pm$ SEM of 3 biological replicates). White and gray shadings represent day and night, respectively.

(I) Hypocotyl length measurements from the indicated lines grown for 14 days in SDs (in gray, mean $\pm$ SEM, n = 16-22).

(A-I) *p < 0.05, **p < 0.01, ***p < 0.001 Student's t test (A-H) or Tukey's multiple comparisons test (I). See also Figures S2 and S3.

electrophoretic mobility shift assays (EMSAs) with proteins expressed in an *in vitro* transcription and translation system (Martínez-García et al., 2000). In these experiments, the binding of FLAG-PIF3 to a *PIL1* promoter fragment containing two G-boxes was reduced in the presence of HA-GI (Figure S2H), even though FLAG-PIF3 levels remained unchanged and GI does not bind the DNA probe directly (Figures S2I and S2J). A similar result was also obtained for FLAG-PIF5 (Figure S2K). Altogether, consistent

results across this set of *in planta*, *in vivo*, and *in vitro* experiments strongly suggest that GI interaction with PIF proteins directly interferes with their stability as well as with their DNA-binding activity *in vivo*.

In order to characterize the effect of GI on PIF association to genomic targets *in vivo*, we performed chromatin immunoprecipitation (ChIP) assays. To uncouple transcriptional regulation of *PIF* expression by GI, we took advantage of the HA-PIF5

overexpression lines (Lorrain et al., 2008), both in Col-0 and a Glox (GI-YPET-FLAG) background. To further uncouple the effect of GI on PIF stability, we also leveraged the PIF3-ECFP-HA overexpression line because a detailed characterization of this line indicated that the ECFP tag might increase PIF3 stability (Janczak et al., 2015) without affecting its primary functions such as DNA binding and phyB-mediated degradation. Specifically, we observed that the accumulation of the PIF3-ECFP-HA fusion protein showed no significant differences when GI is overexpressed both in tobacco and in *Arabidopsis* (Figures S3A–S3C). Furthermore, a treatment with paclobutrazol (PAC), previously shown to affect PIF3 stability (Li et al., 2016), was also ineffective in promoting the degradation of PIF3-ECFP-HA as opposed to PIF5-HA (Figures S3D and S3E), further suggesting that the ECFP tag may be stabilizing it. Despite this increased stability, the fusion protein is still sensitive to degradation through major regulatory mechanisms such as phyB-mediated degradation, and when expressed under the control of its endogenous promoter, it accumulates during the night period as expected (Soy et al., 2012) (Figure S3G). Importantly, it also proved to be functional because plants overexpressing it display longer hypocotyls under different light conditions (Kim et al., 2003) and show upregulation of PIF target genes such as *PIL1* (Figure S3F). Using these lines in our ChIP experiments, for both PIF5 and PIF3 we observed a significantly reduced enrichment of the G-box-containing regions in the promoter of *PIL1* in the immunoprecipitated fractions in the presence of GI at all ZTs tested (Figures 2D and S3H) with a subsequent reduction in *PIL1* mRNA expression (Figure 2E). Conversely, mutation of *GI* resulted in a greater enrichment of the same genomic regions when PIF3 was immunoprecipitated (Figures 2F and S3I), and this greater binding resulted in higher *PIL1* expression at night (Figure 2G). Because in the PIF3-ECFP-HA lines PIF3 protein levels are not affected by GI (Figures S3A–S3G), our results suggest that GI affects PIF binding to target chromatin not only indirectly by affecting their stability but also directly by preventing their binding to specific sites within the chromatin.

If differential stability and activity of PIF proteins underlies the increased expression levels of PIF target genes observed in *gi-2* mutant plants, loss of *PIFs* is expected to reverse this phenotype. As predicted by our hypothesis, loss of *PIF3*, *PIF4*, and *PIF5* drastically reduced the expression of *PIL1* and *LONG HYPOCOTYL IN FAR-RED 1 (HFR1)* in *gi-2* seedlings grown under SD conditions (Figures 2H, S3J, and S3K), and this molecular phenotype also results in a reduction of hypocotyl elongation (Figure 2I). Notably, we observed that genes repressed by PIFs such as *PHYTOENE SYNTHASE (PSY)* are downregulated in *gi-2*, and this downregulation was also rescued by the loss of *PIF3*, *PIF4*, and *PIF5* (Figure S3L). These observations suggest that GI-mediated modulation of PIF activity might be relevant for the regulation of multiple PIF regulated processes, including growth and chlorophyll and carotenoid biosynthesis.

GI and PIFs Occupy the Same Targets Genome Wide in a Phase-Dependent Pattern

Even though GI does not seem to bind DNA directly (Figure S2J), previous reports from our laboratory and others have shown that it is able to associate with chromatin (Kim et al., 2013b; Sawa and Kay, 2011; Sawa et al., 2007). We therefore decided to further

investigate this association and to test whether GI may additionally prevent PIF binding to chromatin by occupying common target sites *in vivo*. To this end, we analyzed GI occupancy at the promoter regions bound by PIFs in *Arabidopsis* transgenic lines expressing tagged GI driven by an endogenous promoter fragment. We observed that GI occupies PIF target regions at dusk and more significantly during the early night (ZT8 and 10) but not later in the night (ZT16) (Figure 3A). This trend correlates well with the observed GI accumulation pattern: under SD conditions, *GI* mRNA peaks at around ZT8, with GI protein levels being highest during the early night and then progressively declining (David et al., 2006; Kim et al., 2007; Sawa et al., 2007) (Figure S4A). On the other hand, PIF accumulation increases as the night period progresses (Figure S4A) and, consistently, PIF3 binding to target sites increases following the same trend (Soy et al., 2016) (Figure 3B). This phased protein accumulation and chromatin occupancy pattern of GI and PIFs, together with the effect of GI misregulation on PIF target gene expression, suggests a function of GI in restricting PIF activity during the early night in order to prevent rapid PIF-mediated regulation of gene expression immediately after transitioning to darkness. In support of this hypothesis, interaction between GI and PIFs was observed to increase as the night progresses peaking at ZT15–16 (Figures S4B and S4C), and we observe a more significant effect of GI on PIF binding to target genes at night (Figures 2D–2G, S3H, and S3I), as well as a higher and more rapid induction of PIF target genes upon darkness in the absence of GI compared to wild-type plants (Figure 3C).

To further test our model, we performed genome-wide analyses of GI genomic targets by ChIP followed by deep sequencing (ChIP-seq). These studies revealed that GI preferentially associates to promoter and intergenic regions (hypergeometric test p value = $8.6e-644$ and $6.1e-109$, respectively) (Figures 3D, S4D, and S4E), and its targets include over 35% of genes found to be misregulated in *gi-2* seedlings in a previous study (Kim et al., 2012) (Figure 3E), which is indicative of a function of GI in the regulation of gene expression. Importantly, motif overrepresentation analysis of GI binding sites revealed the G-box as the most highly enriched *cis* element (Figure 3F). Comparison of GI and PIF3 ChIP-seq experiments under SD conditions (performed at ZT8 and ZT16, respectively) revealed that GI binds a significant set of PIF3 targets (p value < 0.01) (Figures 3G, S4F, and S4G), which are enriched in GO terms related to multiple pathways known to be regulated by PIFs (Leivar and Monte, 2014; Leivar and Quail, 2011) (Figure S4H). Moreover, genes shared by GI and PIF3 tend to be expressed at the end of the night and during the morning/early afternoon in SDs, consistent with genes that are induced and repressed by PIFs (Figure S4I). These results suggest a function of GI as a regulator of PIF transcriptional targets genome wide. Further comparison of GI and PIF3 ChIP-seq peaks revealed that both proteins co-localize and bind to the same sites genome wide (Figures 3H, 3I, and S4J), strengthening the hypothesis that the GI binding may affect PIF access to genomic regions. To gain further insight into how GI regulates PIF3 chromatin accessibility, we performed PIF3 quantitative ChIP-seq experiments with internal spike-in standards in the wild-type and *gi-2* backgrounds. As expected, we observed an overall increment of PIF3 binding to chromatin in the absence of *GI* (Figure 3J). Furthermore,




Figure 3. GI and PIFs Occupy the Same Genomic Targets in a Phase-Dependent Pattern

(A and B) GI (A) and PIF3 (B) ChIP assays of 10-day-old seedlings grown in SD conditions and harvested at the indicated ZTs. The enrichment of the specified regions in the immunoprecipitated samples was quantified by qPCR. Values represent mean $\pm$ SEM of 3 independent experiments ($n = 2-4$; * $p < 0.05$, ** $p < 0.01$ Student's t test). To account for experimental noise, the enrichment in each replicate was calculated as % of input and normalized to the enrichment of *pPIL1* fragments in Col-0.

(C) Relative expression of the indicated PIF targets in wild-type (Col-0) and *gi-2* during early night in SDs (mean $\pm$ SEM of 3 biological replicates; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ Student's t test; data from Figures 2A and S2A are re-plotted). White and gray shadings represent day and night, respectively.

(D) Genomic annotation of GI ChIP-seq peaks. The midpoint of the peaks was used for this analysis.

(E) Overlap between GI ChIP-seq targets and *gi-2* DEGs.

(F) Over-represented *cis* elements around the summit of GI ChIP-seq peaks (± 100 -bp flanking region).

(G) Overlap between GI and PIF3 ChIP-seq peaks (hypergeometric test $p < 0.01$).

(H) Metaplot of the signal from PIF3 and GI ChIP-seq plotted over the centers of PIF3 peaks.

(I) Visualization of PIF3 and GI ChIP-seq data in the genomic region encompassing the *PIL1* locus.

(J) Quantitative analysis of the signal in PIF3-EH;Col-0 and PIF3-EH;*gi-2* ChIP-seq plotted over the centers of PIF3 peaks ($n = 7,962$). The ChIP-seq signal in each sample was additionally normalized to a spiked in control (unmethylated lambda DNA).

(legend continued on next page)

genomic regions with the highest GI-binding signal (top 10%) showed a statistically significant greater increment of PIF3 binding in the absence of GI than genomic regions with moderate or weak GI binding (Figures 3K and S4K). Altogether, our results suggest that GI globally affects PIF3 association to chromatin and more specifically regulates a subset of PIF3 target genes by binding to them and hindering PIF access to regulatory regions. This provides an additional timing mechanism to delay PIF activity and adequately phase the expression of PIF targets to the most appropriate time of the day.

The physiological relevance of our findings for output rhythmicity is evidenced by GI function in the regulation of photoperiodic growth, for example, where GI is required to appropriately phase growth rhythms toward the end of the night under SD photoperiods. Growth rhythm measurements in SDs showed that *gi-2* mutant plants display a general growth de-repression, which is especially notable during the early night (Figure S4L), as previously reported (Nozue et al., 2007). Loss of *PIF3*, however, not only partially rescued the average growth rate phenotype in this mutant but also shifted the maximal growth phase to peak later (Figure S4L).

GI Modulation of Light Signaling Affects Circadian Rhythms

The molecular and physiological interactions between GI and PIFs revealed by our studies provide compelling evidence for how the circadian clock and light signaling pathways intersect for the regulation of output pathways. The observation that PIF3 and GI share targets related to circadian rhythms (Figure S4H) and the fact that both proteins bind to the promoter of the core clock gene *CCA1* around the same region *in vivo* (Figure 4A) prompted us to further investigate the relevance of the GI-PIF interaction for the regulation of circadian rhythmicity. Gene-expression analyses implied that PIFs repress the expression of *CCA1* under SD conditions (see expression levels in the quadruple *pifQ* mutant as compared to wild-type plants, Figure 4B). This observation was further confirmed by transient activation assays using the *pCCA1::LUC* reporter, which showed that PIFs are able to repress *pCCA1* activity (Figure S5A). Notably, mutation of *PIFs* alleviated the very low level of *CCA1* expression in *gi-2* mutants (Figures 4B, 4C, and S5B), suggesting that excessive PIF activity in *gi-2* may (at least partially) be responsible for the low amplitude of *CCA1* in this mutant. At the level of *pCCA1* promoter activation, it also proved to be de-repressed in *pif4-101;pif5-1* mutants, more significantly during the dark to light transitions (Figure S5C). Consistent with the gene-expression analyses, loss-of-function mutation of *PIF3*, *PIF4*, and *PIF5* significantly rescued the low amplitude phenotype of *pCCA1* oscillations in *gi-2*, especially in anticipation of the light phase (Figure 4D).

Analysis of *pCCA1* promoter activity in *Arabidopsis* seedlings under free-running conditions revealed that, although consistent with previous reports, *pif3-1* mutation does not show an obvious period phenotype (Viczián et al., 2005), *pif4-101;pif5-1* double

mutants display a lengthened period in our experimental conditions (Figures S5D and S5E). Conversely, PIF overexpression lines displayed shorter periods than wild-type plants (Figures S5F and S5G), confirming the role of PIF proteins in determining the pace of the clock under these conditions. More importantly, loss of *PIF3* in the *gi-2* background increased the amplitude of *pCCA1* oscillations and lengthened the short period of *gi-2* mutants by 1 h (Figures 4E, 4F, S5H, and S5I). The double *gi-2;pif3-1* mutants also maintained *pCCA1* rhythmicity for longer under free-running conditions than *gi-2* seedlings, which displayed less robust oscillations with higher relative amplitude error (RAE) and became arrhythmic after the third day in constant light (Figure 4E). On the other hand, loss of both *PIF4* and *PIF5* also lengthened the period of *gi-2* mutant seedlings by 1 h, but in this case the oscillations in the triple *gi-2;pif4-101;pif5-1* mutants were less robust and seedlings lost rhythmicity very soon under free-running conditions (Figures 4G, 4H, S5J, and S5K). These results not only suggest that PIF misregulation contributes to the circadian phenotype observed in *gi-2* mutants but also that PIF3 and PIF4/PIF5 play partially non-overlapping roles in this process. We subsequently analyzed the role of PIFs in light-responsive expression of *CCA1* by monitoring *pCCA1* activity in 3-day-old etiolated seedlings transferred to light. In these experiments, we observed a significantly higher increase in the light-induced *pCCA1* activation in *pif4-101;pif5-1* compared to wild-type plants (Figure 4I). On the contrary, this activation was strongly repressed in PIF overexpression lines (Figure S5L). In line with our previous results, *PIF* mutation significantly increased *pCCA1* light responsiveness in *gi-2* (Figure 4I). Finally, the role of PIFs in the light input pathway to the oscillator was further investigated by analyzing *pCCA1::LUC* phase resetting in response to light pulses at specific ZTs. These experiments revealed that while the *pif3-1* mutation does not have a significant effect consistent with previous reports (Viczián et al., 2005), *pif4-101;pif5-1* double mutants display a more pronounced phase delay in response to light, especially during the early night (Figure S5M).

Taken together, our results uncover a role of PIF proteins in light input to the circadian system, directly linking light signaling to *CCA1* transcriptional regulation. We also show that modulation of PIF activity by GI is not only required to adequately phase output rhythms such as growth but also for proper clock progression and light input.

DISCUSSION

Precise timing of physiological processes by the circadian clock provides an adaptive advantage for most organisms (Millar, 2016). In plants, many developmental processes, including photoperiodic growth, are tightly coordinated by the circadian clock, which integrates a variety of internal and external signals (Greenham and McClung, 2015; Sanchez and Kay, 2016). The molecular features required for the efficient decoding and integration of the multiple stimuli perceived within these complex

(K) Boxplot of the signal difference between quantitative PIF3-EH;Col-0 and PIF3-EH;*gi-2* ChIP-seq over GI peaks ranked by increasing signal and divided into 10 groups (deciles 1–10). Statistically significant differences between mean values by Welch's two-sample t test in each decile relative to the 10th decile are shown (**p < 0.01).

See also Figure S4 and Table S1.




Figure 4. GI Modulation of Light Signaling Affects Circadian Rhythms

(A) Visualization of PIF3 and GI ChIP-seq data in the genomic region encompassing the *CCA1* locus.

(B and C) Relative expression of *CCA1* in the indicated backgrounds at ZT24 in SDs (mean $\pm$ SEM of 3 biological replicates).

(D) Bioluminescence analysis of *pCCA1::LUC* in seedlings in SD conditions (mean $\pm$ SEM, $n = 12$). White and gray bars represent day and night, respectively. (E and G) Bioluminescence analysis of *pCCA1::LUC* in the indicated backgrounds in constant light (LL) (mean $\pm$ SEM, $n = 12$). Plants were entrained in SDs for 7 days.

(F and H) Period length estimations of *pCCA1::LUC* reporter expression in the experiments shown in (E) and (G), respectively, as analyzed by fast Fourier transformed non-linear least squares (FFT-NLLS) ($n = 12$).

(I) Bioluminescence analysis of *pCCA1::LUC* activation in 3-day-old etiolated seedlings transferred to light (mean $\pm$ SEM, $n = 12$).

(B–I) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ Student's *t* test. See also Figure S5.

regulatory networks are only just becoming apparent. Here, we provide evidence for a mechanism directly linking a component of the circadian oscillator to the gating of light signaling pathways (Figure 5). Our evidence supports a model where GI directly regulates the activity of the master regulators of light signaling in plants, the PIF proteins, thereby gating the sensitivity to external signals to ultimately shape plant development across the light-dark cycle.

GI modulates light signaling by globally interfering with PIF activity. Importantly, our data indicate that GI affects PIF activity at multiple levels, including *PIF* transcription, PIF degradation, and PIF access to target chromatin. Our observation that GI promotes PIF phosphorylation is especially intriguing because phosphorylation of PIF proteins, which leads to their subsequent ubiquitination and degradation, is a key module of the signal transduction pathway by which photoactivated photoreceptors trigger the expression of light-responsive genes (Ni et al.,

2013; Xu et al., 2015). GI interacts with phyB in yeast and *in planta* (Yeom et al., 2014), but it is unknown whether this interaction occurs *in vivo* and what its functional implications may be. In parallel, a recent study has shown that photoregulatory protein kinases (PPKs) catalyze the photoactivated-phy-induced phosphorylation of PIF3 (Ni et al., 2017). These kinases interact with the Evening Complex through phyB and are involved in circadian rhythmicity and hypocotyl elongation (Zheng et al., 2018); more recently, their function in the latter has been shown to involve synergistic and antagonistic interactions with *CCA1* and *RGA*, respectively (Zheng et al., 2018). It is tempting to speculate that GI may enhance interaction of PIFs with phyB and/or specific protein kinases, such as PPKs. Detailed studies will be required to unravel how GI mediates phosphorylation of PIFs, if it is involved in the aforementioned pathways, what other co-factors are required, and its functional relevance in the context of light signal transduction.




Figure 5. Model Depicting the Function of GI in the Circadian Gating of Light Signaling Pathways

As GI accumulates during the early night, it prevents the binding of the PIFs to chromatin through direct interaction and competition for target promoter regions. Progressive degradation of GI later at night enables the release of the PIFs, which accumulate and access target promoter regions at this time. The proposed mechanism is required not only to adequately phase output rhythms such as growth but also to set the pace of the clock.

The finding that GI affects PIF binding to target promoters not only through direct binding but also by occupying common target regions is a remarkable one and provides evidence of a mechanism that differentiates GI from the other core clock components, such as EARLY FLOWERING 3 (ELF3) and the PSEUDO-RESPONSE REGULATORS (PRRs), which interact with PIFs and repress their activity at growth-promoting genes without affecting their stability and/or their association to target regions (Martín et al., 2018; Nieto et al., 2015; Soy et al., 2016; Zhu et al., 2016). Notably, our genome-wide studies extend GI function as a general regulator of PIFs well beyond the control of hypocotyl elongation, affecting a wide array of pathways in which these transcriptional hubs are involved. To what extent and how GI regulates these other transcriptional networks, what timing properties are delivered by it, and what its impact is on overall plant fitness will be important questions to be addressed in future studies.

The biochemical function of GI, a key component of the circadian oscillator in plants, has long been elusive, partially due to the lack of well-characterized functional domains within the protein and the broad array of mutant allele phenotypes that it displays. The notion of GI as a scaffold protein is widely accepted, given its ability to interact with a multiplicity of partners (Mishra and Panigrahi, 2015). More recently, a function as a holdase that enhances HSP90/HSP70-mediated maturation of the blue-light photoreceptor ZEITLUPE (ZTL) (Cha et al., 2017) has been proposed. Our work takes an important step toward more broadly understanding the molecular function of GI and how it is able to exert its regulatory functions across a wide swath of physiology. We find that GI binds the promoter regions of thousands of genes involved in diverse biological processes. Even

though experiments were performed under different conditions, over a third of genes found to be misregulated in *gi-2* in a previous study are indeed bound by GI, suggesting direct regulation of GI target genes at the transcriptional level. Given that GI does not appear to associate to the DNA directly, an important question will be to uncover the mechanisms by which it is recruited to the chromatin. The finding of the G-box as the most highly enriched motif around GI ChIP-seq peaks points to a G-box-binding transcription factor (or different ones) involved in this recruitment. A plausible candidate would be ELONGATED HYPOCOTYL 5 (HY5). HY5 plays an antagonistic role to PIFs and binds a common set of target promoters in a phased pattern (Toledo-Ortiz et al., 2014), similar to GI. Nevertheless, it is very likely that other DNA-binding transcription factors are also involved because other motifs were also overrepresented around GI peaks and GI is recruited to an ample set of genes that are not bound by PIFs. Whether GI is targeted through the recognition of specific chromatin features such as specific histone marks is also unknown. The identification of the factors that recruit GI to the chromatin, as well as those that are recruited by GI, will be necessary to fully understand the role of GI in the regulation of transcription.

Finally, our study provides evidence on the involvement of PIFs in the light input pathway to the oscillator and how GI modulation of PIF activity is required not only to regulate output processes but also to maintain proper clock progression. Earlier work suggested a role for PIF proteins in the circadian clock (Martínez-García et al., 2000), but this implication could not be subsequently confirmed using PIF3 single mutants and overexpression lines (Viczián et al., 2005) likely because of the high level of redundancy among PIFs. More recently, ChIP-seq experiments have revealed the binding of PIFs to the promoters of clock genes *in vivo* (Oh et al., 2012; Zhang et al., 2013), and the interaction between PIFs and several clock components has been reported (Martín et al., 2018; Nieto et al., 2015; Soy et al., 2016; Zhu et al., 2016). Additionally, a recent study has shown the implication of PIFs in metabolic signaling to the clock (Shor et al., 2017). Our results show that PIFs can directly repress *CCA1* during the night, most remarkably at the dark-to-light transitions, and PIF function is required for period length determination. Successive mutation of *PIFs* in *gi-2* progressively increased *CCA1* amplitude in this mutant and alleviated its short period phenotype, which suggests that excessive PIF activity is at least partially responsible for the canonical clock phenotype of *gi-2* mutants. Nevertheless, mutation of *PIF* genes only partially rescued *gi-2* circadian phenotypes, implicating that additional mechanisms take place in the regulation of *CCA1* expression by GI and in GI function in the circadian system. Different PIFs also seemed to affect *gi-2* rhythmicity differently, which suggests that they may have both overlapping and distinct roles in the system. This could provide mechanisms for differential transcriptional regulation of, or physical interaction with, other clock components. It is also remarkable that the effect of *PIF4* and *PIF5* mutation on *CCA1* phase resetting in response to light is more pronounced in the middle of the night. The specific function of the different PIFs in the circadian system, their dynamics in time, and their interaction with other input pathways will need to be unraveled to fully understand how clock phase is set in response to light.

Altogether, our study assigns a role to GI as a pivotal integrator of external and internal signals to regulate both the circadian clock and its output in the context of transcriptional regulation and provides a mechanistic framework to further investigate the circadian gating of plant development and how light signals are transmitted into the circadian system.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **CONTACT FOR REAGENT AND RESOURCE SHARING**
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Plant Material and Growth Conditions
- **METHOD DETAILS**
 - Construction of Binary Vectors
 - Yeast Two-Hybrid Analyses
 - *In Vitro* Pull-Downs
 - Transient Expression in *N. benthamiana*
 - Protein Immunoprecipitation
 - Protein Stability and Accumulation Analyses
 - Western Blot Detection and Quantitation
 - Electrophoretic Mobility Shift Assay (EMSA)
 - RNA Extraction and qRT-PCR
 - Transactivation Assays
 - Chromatin Immunoprecipitation (ChIP)
 - ChIP-seq
 - Physiological Measurements
 - Clock Activity Measurements
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - Mean Comparisons
 - ChIP-seq Data Analysis
 - Genome-Wide Gene Expression Analyses
 - Growth Rate Calculations
- **DATA AND SOFTWARE AVAILABILITY**

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.devcel.2019.04.030>.

ACKNOWLEDGMENTS

We thank R. McClung and J. Chory for seeds and reagents and N. Buceta, M. Akhter, and S. King for technical assistance. We also thank J. Gallego-Bartolome, S. Sanchez, D. Alabadi, and D. Nusinow for critical reading of the manuscript. This work was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award numbers RO1GM067837 and RO1GM056006 to S.A.K. J.N.M. and K.N. acknowledge United States Department of Agriculture USDA NIFA project CA-D-PLB-7226-H and UC Davis for funding. S.E.J. is an investigator from the Howard Hughes Medical Institute.

AUTHOR CONTRIBUTIONS

Conceptualization, M.A.N. and S.A.K.; Investigation, M.A.N., W.L., T.D., M.S., J.L.P.-P., and K.N.; Resources, J.N.M., S.E.J., and S.A.K.; Writing, M.A.N. and S.A.K.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: September 2, 2018

Revised: February 16, 2019

Accepted: April 17, 2019

Published: May 16, 2019

REFERENCES

- Alabadi, D., Oyama, T., Yanovsky, M.J., Harmon, F.G., Más, P., and Kay, S.A. (2001). Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. *Science* 293, 880–883.
- Berns, M.C., Nordström, K., Cremer, F., Tóth, R., Hartke, M., Simon, S., Klasen, J.R., Büchel, I., and Coupland, G. (2014). Evening expression of Arabidopsis GIGANTEA is controlled by combinatorial interactions among evolutionarily conserved regulatory motifs. *Plant Cell* 26, 3999–4018.
- Cao, S., Ye, M., and Jiang, S. (2005). Involvement of GIGANTEA gene in the regulation of the cold stress response in Arabidopsis. *Plant Cell Rep.* 24, 683–690.
- Castillon, A., Shen, H., and Huq, E. (2007). Phytochrome interacting factors: central players in phytochrome-mediated light signaling networks. *Trends Plant Sci.* 12, 514–521.
- Cha, J.Y., Kim, J., Kim, T.S., Zeng, Q., Wang, L., Lee, S.Y., Kim, W.Y., and Somers, D.E. (2017). GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE in the Arabidopsis circadian clock. *Nat. Commun.* 8, 3.
- Covington, M.F., Maloof, J.N., Straume, M., Kay, S.A., and Harmer, S.L. (2008). Global transcriptome analysis reveals circadian regulation of key pathways in plant growth and development. *Genome Biol.* 9, R130.
- David, K.M., Armbruster, U., Tama, N., and Putterill, J. (2006). Arabidopsis GIGANTEA protein is post-transcriptionally regulated by light and dark. *FEBS Lett.* 580, 1193–1197.
- Davière, J.M., and Achard, P. (2016). A pivotal role of DELLAs in regulating multiple hormone signals. *Mol. Plant* 9, 10–20.
- de Lucas, M., Davière, J.M., Rodríguez-Falcón, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S., Fankhauser, C., Blázquez, M.A., Titarenko, E., and Prat, S. (2008). A molecular framework for light and gibberellin control of cell elongation. *Nature* 451, 480–484.
- de Montaigu, A., Giakountis, A., Rubin, M., Tóth, R., Cremer, F., Sokolova, V., Porri, A., Reymond, M., Weinig, C., and Coupland, G. (2015). Natural diversity in daily rhythms of gene expression contributes to phenotypic variation. *Proc. Natl. Acad. Sci. USA* 112, 905–910.
- Earley, K.W., Haag, J.R., Pontes, O., Opper, K., Juehne, T., Song, K., and Pikaard, C.S. (2006). Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J.* 45, 616–629.
- Feng, S., Martinez, C., Gusmaroli, G., Wang, Y., Zhou, J., Wang, F., Chen, L., Yu, L., Iglesias-Pedraz, J.M., Kircher, S., et al. (2008). Coordinated regulation of Arabidopsis thaliana development by light and gibberellins. *Nature* 451, 475–479.
- Fornara, F., de Montaigu, A., Sánchez-Villalreal, A., Takahashi, Y., Ver Loren van Themaat, E., Huettel, B., Davis, S.J., and Coupland, G. (2015). The GI-CDF module of Arabidopsis affects freezing tolerance and growth as well as flowering. *Plant J.* 81, 695–706.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G., and Putterill, J. (1999). Gigantea: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. *EMBO J.* 18, 4679–4688.
- Fujimori, T., Yamashino, T., Kato, T., and Mizuno, T. (2004). Circadian-controlled basic/helix-loop-helix factor, PIL6, implicated in light-signal transduction in Arabidopsis thaliana. *Plant Cell Physiol.* 45, 1078–1086.
- Gould, P.D., Locke, J.C., Larue, C., Southern, M.M., Davis, S.J., Hanano, S., Moyle, R., Milich, R., Putterill, J., Millar, A.J., et al. (2006). The molecular basis

- of temperature compensation in the Arabidopsis circadian clock. *Plant Cell* 18, 1177–1187.
- Greenham, K., and McClung, C.R. (2015). Integrating circadian dynamics with physiological processes in plants. *Nat. Rev. Genet.* 16, 598–610.
- Harmer, S.L., Hogenesch, J.B., Straume, M., Chang, H.S., Han, B., Zhu, T., Wang, X., Kreps, J.A., and Kay, S.A. (2000). Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. *Science* 290, 2110–2113.
- Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* 38, 576–589.
- Hellens, R.P., Allan, A.C., Friel, E.N., Bolitho, K., Grafton, K., Templeton, M.D., Karunairatnam, S., Gleave, A.P., and Laing, W.A. (2005). Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants. *Plant Methods* 1, 13.
- Hornitschek, P., Lorrain, S., Zoete, V., Michielin, O., and Fankhauser, C. (2009). Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO J.* 28, 3893–3902.
- Huq, E., and Quail, P.H. (2002). PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in Arabidopsis. *EMBO J.* 21, 2441–2450.
- Huq, E., Tepperman, J.M., and Quail, P.H. (2000). GIGANTEA is a nuclear protein involved in phytochrome signaling in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 97, 9789–9794.
- Janczak, M., Bukowski, M., Górecki, A., Dubin, G., Dubin, A., and Wladyka, B. (2015). A systematic investigation of the stability of green fluorescent protein fusion proteins. *Acta Biochim. Pol.* 62, 407–411.
- Johnson, L.M., Du, J., Hale, C.J., Bischof, S., Feng, S., Chodavarapu, R.K., Zhong, X., Marson, G., Pellegrini, M., Segal, D.J., et al. (2014). SRA- and SET-domain-containing proteins link RNA polymerase V occupancy to DNA methylation. *Nature* 507, 124–128.
- Kamioka, M., Takao, S., Suzuki, T., Taki, K., Higashiyama, T., Kinoshita, T., and Nakamichi, N. (2016). Direct repression of evening genes by CIRCADIAN CLOCK-ASSOCIATED1 in the Arabidopsis circadian clock. *Plant Cell* 28, 696–711.
- Kim, J., Yi, H., Choi, G., Shin, B., Song, P.S., and Choi, G. (2003). Functional characterization of phytochrome interacting factor 3 in phytochrome-mediated light signal transduction. *Plant Cell* 15, 2399–2407.
- Kim, W.Y., Ali, Z., Park, H.J., Park, S.J., Cha, J.Y., Perez-Hormaeche, J., Quintero, F.J., Shin, G., Kim, M.R., Qiang, Z., et al. (2013a). Release of SOS2 kinase from sequestration with GIGANTEA determines salt tolerance in Arabidopsis. *Nat. Commun.* 4, 1352.
- Kim, W.Y., Fujiwara, S., Suh, S.S., Kim, J., Kim, Y., Han, L., David, K., Putterill, J., Nam, H.G., and Somers, D.E. (2007). ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature* 449, 356–360.
- Kim, Y., Lim, J., Yeom, M., Kim, H., Kim, J., Wang, L., Kim, W.Y., Somers, D.E., and Nam, H.G. (2013b). ELF4 regulates GIGANTEA chromatin access through subnuclear sequestration. *Cell Rep.* 3, 671–677.
- Kim, Y., Yeom, M., Kim, H., Lim, J., Koo, H.J., Hwang, D., Somers, D., and Nam, H.G. (2012). GIGANTEA and EARLY FLOWERING 4 in Arabidopsis exhibit differential phase-specific genetic influences over a diurnal cycle. *Mol. Plant* 5, 678–687.
- Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25.
- Legris, M., Nieto, C., Sellaro, R., Prat, S., and Casal, J.J. (2017). Perception and signalling of light and temperature cues in plants. *Plant J.* 90, 683–697.
- Leivar, P., and Monte, E. (2014). PIFs: systems integrators in plant development. *Plant Cell* 26, 56–78.
- Leivar, P., Monte, E., Oka, Y., Liu, T., Carle, C., Castillon, A., Huq, E., and Quail, P.H. (2008). Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr. Biol.* 18, 1815–1823.
- Leivar, P., and Quail, P.H. (2011). PIFs: pivotal components in a cellular signaling hub. *Trends Plant Sci.* 16, 19–28.
- Li, K., Yu, R., Fan, L.M., Wei, N., Chen, H., and Deng, X.W. (2016). DELLA-mediated PIF degradation contributes to coordination of light and gibberellin signalling in Arabidopsis. *Nat. Commun.* 7, 11868.
- Locke, J.C., Kozma-Bognár, L., Gould, P.D., Fehér, B., Kevei, E., Nagy, F., Turner, M.S., Hall, A., and Millar, A.J. (2006). Experimental validation of a predicted feedback loop in the multi-oscillator clock of Arabidopsis thaliana. *Mol. Syst. Biol.* 2, 59.
- Lorrain, S., Allen, T., Duek, P.D., Whitelam, G.C., and Fankhauser, C. (2008). Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J.* 53, 312–323.
- Martín, G., Rovira, A., Veciana, N., Soy, J., Toledo-Ortiz, G., Gommers, C.M.M., Boix, M., Henriques, R., Minguet, E.G., Alabadi, D., et al. (2018). Circadian waves of transcriptional repression shape PIF-regulated photoperiod-responsive growth in Arabidopsis. *Curr. Biol.* 28, 311–318.e5.
- Martínez-García, J.F., Huq, E., and Quail, P.H. (2000). Direct targeting of light signals to a promoter element-bound transcription factor. *Science* 288, 859–863.
- Martin-Tryon, E.L., Kreps, J.A., and Harmer, S.L. (2007). GIGANTEA acts in blue light signaling and has biochemically separable roles in circadian clock and flowering time regulation. *Plant Physiol.* 143, 473–486.
- Millar, A.J. (2016). The intracellular dynamics of circadian clocks reach for the light of ecology and evolution. *Annu. Rev. Plant Biol.* 67, 595–618.
- Mishra, P., and Panigrahi, K.C. (2015). GIGANTEA - an emerging story. *Front. Plant Sci.* 6, 8.
- Mizoguchi, T., Wheatley, K., Hanzawa, Y., Wright, L., Mizoguchi, M., Song, H.R., Carré, I.A., and Coupland, G. (2002). LHY and CCA1 are partially redundant genes required to maintain circadian rhythms in Arabidopsis. *Dev. Cell* 2, 629–641.
- Mizoguchi, T., Wright, L., Fujiwara, S., Cremer, F., Lee, K., Onouchi, H., Mouradov, A., Fowler, S., Kamada, H., Putterill, J., et al. (2005). Distinct roles of GIGANTEA in promoting flowering and regulating circadian rhythms in Arabidopsis. *Plant Cell* 17, 2255–2270.
- Mockler, T.C., Michael, T.P., Priest, H.D., Shen, R., Sullivan, C.M., Givan, S.A., McEntee, C., Kay, S.A., and Chory, J. (2007). The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. *Cold Spring Harb. Symp. Quant. Biol.* 72, 353–363.
- Möglich, A., Yang, X., Ayers, R.A., and Moffat, K. (2010). Structure and function of plant photoreceptors. *Annu. Rev. Plant Biol.* 61, 21–47.
- Moore, A., Zielinski, T., and Millar, A.J. (2014). Online period estimation and determination of rhythmicity in circadian data, using the BioDare data infrastructure. *Methods Mol. Biol.* 1158, 13–44.
- Ni, W., Xu, S.L., Chalkley, R.J., Pham, T.N., Guan, S., Maltby, D.A., Burlingame, A.L., Wang, Z.Y., and Quail, P.H. (2013). Multisite light-induced phosphorylation of the transcription factor PIF3 is necessary for both its rapid degradation and concomitant negative feedback modulation of photoreceptor phyB levels in Arabidopsis. *Plant Cell* 25, 2679–2698.
- Ni, W., Xu, S.L., González-Grandío, E., Chalkley, R.J., Huhmer, A.F.R., Burlingame, A.L., Wang, Z.Y., and Quail, P.H. (2017). PPKs mediate direct signal transfer from phytochrome photoreceptors to transcription factor PIF3. *Nat. Commun.* 8, 15236.
- Nieto, C., López-Salmerón, V., Davière, J.M., and Prat, S. (2015). ELF3-PIF4 interaction regulates plant growth independently of the Evening Complex. *Curr. Biol.* 25, 187–193.
- Nito, K., Wong, C.C., Yates, J.R., 3rd, and Chory, J. (2013). Tyrosine phosphorylation regulates the activity of phytochrome photoreceptors. *Cell Rep.* 3, 1970–1979.
- Nohales, M.A., and Kay, S.A. (2016). Molecular mechanisms at the core of the plant circadian oscillator. *Nat. Struct. Mol. Biol.* 23, 1061–1069.
- Nozue, K., Covington, M.F., Duek, P.D., Lorrain, S., Fankhauser, C., Harmer, S.L., and Maloof, J.N. (2007). Rhythmic growth explained by coincidence between internal and external cues. *Nature* 448, 358–361.

- Nusinow, D.A., Helfer, A., Hamilton, E.E., King, J.J., Imaizumi, T., Schultz, T.F., Farré, E.M., and Kay, S.A. (2011). The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475, 398–402.
- Oh, E., Zhu, J.Y., and Wang, Z.Y. (2012). Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nat. Cell Biol.* 14, 802–809.
- Oliverio, K.A., Crepy, M., Martin-Tryon, E.L., Milich, R., Harmer, S.L., Putterill, J., Yanovsky, M.J., and Casal, J.J. (2007). GIGANTEA regulates phytochrome A-mediated photomorphogenesis independently of its role in the circadian clock. *Plant Physiol.* 144, 495–502.
- Park, D.H., Somers, D.E., Kim, Y.S., Choy, Y.H., Lim, H.K., Soh, M.S., Kim, H.J., Kay, S.A., and Nam, H.G. (1999). Control of circadian rhythms and photo-periodic flowering by the Arabidopsis GIGANTEA gene. *Science* 285, 1579–1582.
- Pedmale, U.V., Huang, S.C., Zander, M., Cole, B.J., Hetzel, J., Ljung, K., Reis, P.A.B., Sridevi, P., Nito, K., Nery, J.R., et al. (2016). Cryptochromes interact directly with PIFs to control plant growth in limiting blue light. *Cell* 164, 233–245.
- Pruneda-Paz, J.L., Breton, G., Nagel, D.H., Kang, S.E., Bonaldi, K., Doherty, C.J., Ravelo, S., Galli, M., Ecker, J.R., and Kay, S.A. (2014). A genome-scale resource for the functional characterization of Arabidopsis transcription factors. *Cell Rep.* 8, 622–632.
- Pruneda-Paz, J.L., Breton, G., Para, A., and Kay, S.A. (2009). A functional genomics approach reveals CHE as a component of the Arabidopsis circadian clock. *Science* 323, 1481–1485.
- R Development Core Team (2016). R: a language and environment for statistical computing (R Foundation for Statistical Computing). <https://www.R-project.org/>.
- Salomé, P.A., and McClung, C.R. (2005). PSEUDO-RESPONSE REGULATOR 7 and 9 are partially redundant genes essential for the temperature responsiveness of the Arabidopsis circadian clock. *Plant Cell* 17, 791–803.
- Sanchez, S.E., and Kay, S.A. (2016). The plant circadian clock: From a simple timekeeper to a complex developmental manager. *Cold Spring Harb. Perspect. Biol.* 8.
- Sawa, M., and Kay, S.A. (2011). GIGANTEA directly activates Flowering Locus T in Arabidopsis thaliana. *Proc. Natl. Acad. Sci. USA* 108, 11698–11703.
- Sawa, M., Nusinow, D.A., Kay, S.A., and Imaizumi, T. (2007). FKF1 and GIGANTEA complex formation is required for day-length measurement in Arabidopsis. *Science* 318, 261–265.
- Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675.
- Shen, H., Zhu, L., Castillon, A., Majee, M., Downie, B., and Huq, E. (2008). Light-induced phosphorylation and degradation of the negative regulator PHYTOCHROME-INTERACTING FACTOR1 from Arabidopsis depend upon its direct physical interactions with photoactivated phytochromes. *Plant Cell* 20, 1586–1602.
- Shen, L., Shao, N., Liu, X., and Nestler, E. (2014). ngs.plot: quick mining and visualization of next-generation sequencing data by integrating genomic databases. *BMC Genomics* 15, 284.
- Shor, E., Paik, I., Kangisser, S., Green, R., and Huq, E. (2017). PHYTOCHROME INTERACTING FACTORS mediate metabolic control of the circadian system in Arabidopsis. *New Phytol.* 215, 217–228.
- Soy, J., Leivar, P., González-Schain, N., Martín, G., Diaz, C., Sentandreu, M., Al-Sady, B., Quail, P.H., and Monte, E. (2016). Molecular convergence of clock and photosensory pathways through PIF3-TOC1 interaction and co-occupancy of target promoters. *Proc. Natl. Acad. Sci. USA* 113, 4870–4875.
- Soy, J., Leivar, P., González-Schain, N., Sentandreu, M., Prat, S., Quail, P.H., and Monte, E. (2012). Phytochrome-imposed oscillations in PIF3 protein abundance regulate hypocotyl growth under diurnal light/dark conditions in Arabidopsis. *Plant J.* 71, 390–401.
- Suárez-López, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F., and Coupland, G. (2001). CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. *Nature* 410, 1116–1120.
- Thomas, P.D., Campbell, M.J., Kejariwal, A., Mi, H., Karlak, B., Daverman, R., Diemer, K., Muruganujan, A., and Narechania, A. (2003). PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res.* 13, 2129–2141.
- Toledo-Ortiz, G., Johansson, H., Lee, K.P., Bou-Torrent, J., Stewart, K., Steel, G., Rodríguez-Concepción, M., and Halliday, K.J. (2014). The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS Genet.* 10, e1004416.
- Viczián, A., Kircher, S., Fejes, E., Millar, A.J., Schäfer, E., Kozma-Bognár, L., and Nagy, F. (2005). Functional characterization of phytochrome interacting factor 3 for the Arabidopsis thaliana circadian clockwork. *Plant Cell Physiol.* 46, 1591–1602.
- Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis (Springer-Verlag).
- Xu, H., Liu, Q., Yao, T., and Fu, X. (2014). Shedding light on integrative GA signaling. *Curr. Opin. Plant Biol.* 21, 89–95.
- Xu, X., Paik, I., Zhu, L., and Huq, E. (2015). Illuminating progress in phytochrome-mediated light signaling pathways. *Trends Plant Sci.* 20, 641–650.
- Yeom, M., Kim, H., Lim, J., Shin, A.Y., Hong, S., Kim, J.I., and Nam, H.G. (2014). How do phytochromes transmit the light quality information to the circadian clock in Arabidopsis? *Mol. Plant* 7, 1701–1704.
- Zhang, Y., Liu, T., Meyer, C.A., Eickhout, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* 9, R137.
- Zhang, Y., Mayba, O., Pfeiffer, A., Shi, H., Tepperman, J.M., Speed, T.P., and Quail, P.H. (2013). A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in Arabidopsis. *PLoS Genet.* 9, e1003244.
- Zheng, H., Zhang, F., Wang, S., Su, Y., Ji, X., Jiang, P., Chen, R., Hou, S., and Ding, Y. (2018). MLK1 and MLK2 coordinate RGA and CCA1 activity to regulate hypocotyl elongation in Arabidopsis thaliana. *Plant Cell* 30, 67–82.
- Zhu, J.Y., Oh, E., Wang, T., and Wang, Z.Y. (2016). TOC1-PIF4 interaction mediates the circadian gating of thermoresponsive growth in Arabidopsis. *Nat. Commun.* 7, 13692.
- Zielinski, T., Moore, A.M., Troup, E., Halliday, K.J., and Millar, A.J. (2014). Strengths and limitations of period estimation methods for circadian data. *PLoS One* 9, e96462.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-FLAG M2	Sigma	Cat# F1804; RRID:AB_262044
Anti-FLAG-HRP	Sigma	Cat#A8592; RRID:AB_439702
Anti-HA 3F10	Roche	Cat# 11867423001; RRID:AB_2314622
Anti-HA-HRP	Roche	Cat# 12013819001; RRID: AB_390917
Anti-Actin	Millipore	Cat# MAB1501; RRID:AB_2223041
Anti-Mouse-HRP	BioRad	Cat#1706516
Bacterial and Virus Strains		
<i>E. coli</i> ccdB survival	Invitrogen	A10460
<i>E. coli</i> TOP10	Invitrogen	C404003
<i>Agrobacterium tumefaciens</i> strain GV3101	N/A	N/A
<i>Agrobacterium tumefaciens</i> strain C58	N/A	N/A
Chemicals, Peptides, and Recombinant Proteins		
Hygromycin B	Invitrogen	Cat# 10687010
Glufosinate ammonium (BASTA)	Sigma	Cat#45520
PMSF	Sigma	Cat# P7626
Complete EDTA-free Protease Inhibitor	Roche	Cat# 11836170001
Phosphatase Inhibitor 2	Sigma	Cat#P5726
Phosphatase Inhibitor 3	Sigma	Cat#P0044
MG-132	Peptides International	Cat#3175-v
Protein G Dynabeads	Invitrogen	Cat# 10004D
GlycoBlue	Invitrogen	Cat# AM9515
D-Luciferin, potassium salt	Biosynth AG	Cat#L-8220
Critical Commercial Assays		
Gateway BP clonase enzyme mix	Invitrogen	Cat#11789020
Gateway LR clonase enzyme mix	Invitrogen	Cat#11791100
ProQuest Two-Hybrid System	Invitrogen	Cat#PQ1000101
TnT SP6 High-Yield Wheat Germ Protein Expression System	Promega	Cat#L3260
DC Protein Assay	BioRad	Cat#5000116
Supersignal West Pico, Dura, and Femto substrates	Pierce	Cat#34577, 34075, 34095
iScript cDNA synthesis kit	BioRad	Cat#1708890
Maxima SYBR Green qPCR Master Mix	Thermo Scientific	Cat#K0253
Ovation Ultralow V2 kit	NuGEN	Cat# 0344NB-A01
Deposited Data		
ChIP-seq data	This paper	GEO: GSE129865
Experimental Models: Organisms/Strains		
Col-0	N/A	N/A
<i>gi-2</i>	Fowler et al., 1999	N/A
Glox	David et al., 2006	N/A
PIF5-HA	Lorrain et al., 2008	N/A
PIF4-Flash	Pedmale et al., 2016	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<i>pif3-1</i>	Kim et al., 2003	SALK_030753
<i>pif4-101</i>	Lorrain et al., 2008	Garlic_114_G06
<i>pif5-1</i>	Fujimori et al., 2004	SALK_087012
<i>pifQ</i>	Leivar et al., 2008	N/A
CCA1::LUC	Salomé and McClung, 2005	N/A
<i>gi-2;pif3-1</i>	This paper	N/A
<i>gi-2;pif4-101</i>	This paper	N/A
<i>gi-2;pif5-1</i>	This paper	N/A
<i>gi-2;pif4-101;pif5-1</i>	This paper	N/A
GI-YPET-Flag	This paper	N/A
pGI::GI-YPET-Flag	This paper	N/A
PIF3-ECFP-HA	This paper	N/A
pPIF3::PIF3-ECFP-HA	This paper	N/A
PIF3-ECFP-HA;GI-YPET-Flag	This paper	N/A
PIF3-ECFP-HA; <i>gi-2</i>	This paper	N/A
pGI::GI-YPET-Flag;pPIF3::PIF3-ECFP-HA	This paper	N/A
PIF5-HA;GI-YPET-Flag	This paper	N/A
pCCA1::LUC, <i>gi-2;pif3-1</i>	This paper	N/A
pCCA1::LUC; <i>gi-2;pif4-101</i>	This paper	N/A
pCCA1::LUC; <i>gi-2;pif5-1</i>	This paper	N/A
pCCA1::LUC; <i>gi-2;pif4-101;pif5-1</i>	This paper	N/A
pCCA1::LUC; PIF3-ECFP-HA	This paper	N/A
pCCA1::LUC; PIF5-HA	This paper	N/A
pCCA1::LUC; PIF4-Flash	This paper	N/A
Oligonucleotides		
Primers used in this study, see Table S2	N/A	N/A
Recombinant DNA		
pD32-GI	This paper	N/A
pD22-PIF1	Pruneda-Paz et al., 2014	N/A
pD22-PIF3	Pruneda-Paz et al., 2014	N/A
pD22-PIF4	Pruneda-Paz et al., 2014	N/A
pD22-PIF5	Pruneda-Paz et al., 2014	N/A
pTNT-HA	Chory lab	N/A
pTNT-Flag	Chory lab	N/A
pTNT-HA-GI	This paper	N/A
pTNT-Flag-PIF1	This paper	N/A
pTNT-Flag-PIF3	This paper	N/A
pTNT-Flag-PIF4	This paper	N/A
pTNT-Flag-PIF5	This paper	N/A
pTNT-Flag-GFP	This paper	N/A
pTNT-Flag-P3-1	This paper	N/A
pTNT-Flag-P3-2	This paper	N/A
pTNT-Flag-P3-3	This paper	N/A
pTNT-Flag-P3-4	This paper	N/A
pB7m34GW	University of Ghent collection	https://gateway.psb.ugent.be
pB-35S::GI-YPET-Flag	This paper	N/A
pB-pGI::GI-YPET-Flag	This paper	N/A
pH7m34GW	University of Ghent collection	https://gateway.psb.ugent.be
pH-35S::PIF3-ECFP-HA	This paper	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
pH-pPIF3::PIF3-ECFP-HA	This paper	N/A
pEarleyGate201	Earley et al., 2006	N/A
pEarleyGate202	Earley et al., 2006	N/A
pEG201-PIF3	This paper	N/A
pEG201-PIF5	This paper	N/A
pEG201-GFP	This paper	N/A
pEG202-PIF1	This paper	N/A
pEG202-PIF3	This paper	N/A
pEG202-PIF4	This paper	N/A
pEG202-PIF5	This paper	N/A
pEG201-GI	This paper	N/A
pEG201-GFP	This paper	N/A
pGreenII 0800-LUC	Hellens et al., 2005	N/A
pGLUCpPIL1	This paper	N/A
pGLUCpCCA1	This paper	N/A
Software and Algorithms		
VisionWorksLS	UVP, LLC	N/A
NIH ImageJ software	Schneider et al., 2012	https://imagej.nih.gov/ij/
CFX Manager	BioRad	Cat#1845000
LabView	National Instruments	N/A
BioDare2	Moore et al., 2014 ; Zielinski et al., 2014	https://www.biodare2.ed.ac.uk
Bowtie (v1.0.0)	Langmead et al., 2009	http://bowtie-bio.sourceforge.net/index.shtml
MACS2 (v 2.1.1.)	Zhang et al., 2008	https://pypi.org/project/MACS2/
NGSplot (v 2.41.4)	Shen et al., 2014	https://github.com/shenlab-sinai/ngsplot
HOMER	Heinz et al., 2010	http://homer.ucsd.edu/homer/download.html
Phaser	Mockler et al., 2007	http://phaser.mocklerlab.org/
PANTHER	Thomas et al., 2003	http://www.pantherdb.org/
R	R Development Core Team, 2016	http://www.R-project.org/
ggplot2 plotting package	Wickham, 2009	https://cran.r-project.org/web/packages/ggplot2/

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Steve A. Kay (stevekay@usc.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Plant Material and Growth Conditions

Wildtype, mutant, and transgenic lines used in this study were *Arabidopsis thaliana* ecotype Columbia 0 (Col-0). *gi-2* ([Fowler et al., 1999](#)), *Glox* ([David et al., 2006](#)), PIF5-HA ([Lorrain et al., 2008](#)), PIF4-Flash ([Pedmale et al., 2016](#)), *pif3-1* (SALK_030753) ([Kim et al., 2003](#)), *pif4-101* (Garlic_114_G06) ([Lorrain et al., 2008](#)), *pif5-1* (SALK_087012) ([Fujimori et al., 2004](#)), *pifQ* ([Leivar et al., 2008](#)), and CCA1::LUC ([Salomé and McClung, 2005](#)) have been previously described. *pif3-1* and *pifQ* lines were obtained from the Arabidopsis Biological Research Center (ABRC) collection. The *gi-2* mutation is a deletion ($\Delta 670-677$) that introduces a premature stop codon and is considered a null mutation predicted to encode a nonfunctional 114-aa polypeptide rather than the 1173-aa GI protein ([Fowler et al., 1999](#)).

Seeds were chlorine gas sterilized and plated on 0.5x Linsmaier and Skoog medium (LS, Caisson Laboratories) with 0.8% agar (Sigma). After stratification in the dark at 4°C for 3 days, plates were transferred to a Percival incubator (Percival-scientific.com)

set to the indicated light conditions with light supplied at $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ by cool-white fluorescent bulbs and a constant temperature of 22°C.

Generation of Plant Lines

gi-2; pif3-1, *gi-2; pif4-101*, *gi-2; pif5-1* and *gi-2; pif4-101; pif5-1* higher order mutants were generated by genetic crosses between single and double mutants, and homozygous mutant lines were identified in the F2 populations by PCR amplification with primers listed in Table S2.

PIF3-ECFP-HA; GI-YPET-Flag and PIF3-ECFP-HA;*gi-2* lines were obtained by crossing a transgenic line overexpressing PIF3-ECFP-HA (hygromycin resistance) with a transgenic line overexpressing GI-YPET-Flag (BASTA resistant and *gi-2*). F3 populations were scored for hygromycin resistance and BASTA resistance/sensitivity. The presence of the *gi-2* allele in both lines was determined by PCR amplification with primers listed in Table S2. PIF5-HA was crossed with GI-YPET-Flag to obtain PIF5-HA; GI-YPET-Flag. F3 populations were scored for seed coat GFP fluorescence and BASTA resistance. The presence of the *gi-2* allele was determined by PCR amplification with primers listed in Table S2. For co-immunoprecipitation studies, F2 segregating populations derived from these crosses were used.

The transgenic PIF3-ECFP-HA and GI-YPET-Flag lines were generated by *Agrobacterium*-mediated floral dip transformation of Col-0 and *gi-2* plants, respectively. To this purpose, *Agrobacterium tumefaciens* strain GV3101 was transformed with the binary vectors pH-35S::PIF3-ECFP-HA, pH-pPIF3::PIF3-ECFP-HA, pB-35S::GI-YPET-Flag, and pB-pGI::GI-YPET-Flag (described below), respectively. Functionality of the fusion proteins was determined phenotypically. PIF3-ECFP-HA overexpressing plants displayed longer hypocotyls under both SD and continuous light and were hyposensitive to red light. On the contrary, GI-YPET-Flag overexpressing plants displayed shorter hypocotyls under both SD and continuous light. The GI-YPET-Flag transgene also rescued the late flowering phenotype of *gi-2* plants.

To obtain the different reporter lines expressing *pCCA1::LUC*, the mutant lines *gi-2; pif3-1*, *gi-2; pif4-101; pif5-1*, and the PIF overexpressing lines PIF3-ECFP-HA, PIF5-HA, and PIF4-Flash were crossed with the *CCA1::LUC* line (BASTA resistance). *pCCA1::LUC* homozygous lines were identified in the F3 population by scoring BASTA resistance and the presence of the different mutant alleles and transgenes was determined by PCR amplification with primers listed in Table S2.

METHOD DETAILS

Construction of Binary Vectors

The binary vectors used for co-localization studies in *Nicotiana benthamiana* and the generation of transgenic *Arabidopsis* lines were constructed by MultiSite Gateway Three-Fragment Vector Technology (Invitrogen). Full-length cDNAs encoding PIF3 (without the stop codon) were amplified by PCR and cloned into the pENTR/D-TOPO vector (Invitrogen) (primers used are listed in Table S2). pENTR-GI-stop (Sawa and Kay, 2011) and pENTR-PIF5 (Pruneda-Paz et al., 2014) have been described earlier. pENTR-PIF5 was mutagenized by PCR using the primers listed in Table S2 to eliminate the stop codon. In the case of the endogenous *PIF3* and *GI* promoter sequences, 2000 bp upstream of the *PIF3* start codon or 2533 bp upstream of the *GI* start codon (Berns et al., 2014) were amplified by PCR using the primers listed in Table S2 and cloned into the pDONR P4-P1R vector (Invitrogen) by Gateway BP recombination reaction (Invitrogen). The pDONR P4-P1R vector containing the 35S promoter sequence was a gift from Joanne Chory (The SALK Institute, La Jolla, CA, United States of America). The sequences of YPET and ECFP-HA (from pEarleyGate102 (Earley et al., 2006)) were subcloned into the pDONR P2R-P3 vector (primer listed in Table S2). pDONR P2R-P3-YPET was subsequently modified by PCR-based mutagenesis to introduce a Flag tag (primers used are listed in Table S2). Selected promoter, gene, and fluorescent tag combinations were finally cloned by MultiSite Gateway reaction (Invitrogen) into either pH7m34GW (35S::PIF3-ECFP-HA and pPIF3::PIF3-ECFP-HA) or pB7m34GW (35S::GI-YPET-Flag and pB-pGI::GI-YPET-Flag) binary destination vectors from the University of Ghent collection (<https://gateway.psb.ugent.be>).

To perform protein stability and transactivation assays in transient expression in *N. benthamiana*, the cDNAs encoding GI, PIF1, PIF3, PIF4, PIF5, and GFP were amplified by PCR from the pENTR/D-TOPO vector (primers used are listed in Table S2) and subcloned into the pDONR207 vector (Invitrogen) by Gateway BP recombination reaction (Invitrogen). Subsequently, they were transferred to either pEarleyGate201 or pEarleyGate202 (Earley et al., 2006) (pEG) binary destination vectors by Gateway LR recombination reaction (Invitrogen). The sequences introduced into these plasmids contained stop codon. Specifically, the different constructs generated and used were as follow: pEG201-GFP, pEG202-GI, pH-35S::PIF3-ECFP-HA, pEG201-PIF3, and pEG201-PIF5 were used for protein stability analyses, and pEG202-PIF1, pEG202-PIF3, pEG202-PIF4, pEG202-PIF5, pEG201-GI, and pEG201-GFP were used for transient transcriptional activation assays. To prepare the reporter constructs for the transactivation assays, the promoter sequences of *PIL1* (2066 bp upstream of the start codon) and *CCA1* (1140 bp upstream of the start codon) were amplified by PCR from genomic DNA (primers used are listed in Table S2) and cloned into the XhoI and BamHI sites of the pGreenII 0800-LUC vector (Hellens et al., 2005).

Yeast Two-Hybrid Analyses

The ProQuest Two-Hybrid System (Invitrogen) was used to perform yeast two-hybrid (Y2H) analyses. The cDNA encoding full-length GI was transferred from the pENTR/D-TOPO vector (Invitrogen) into the pDEST32 vector by Gateway LR recombination reaction (Invitrogen) to generate the bait plasmid. The pDEST22 prey plasmids containing the sequences encoding PIF1, PIF3, PIF4, and PIF5 have been previously described (Pruneda-Paz et al., 2014). Empty pDEST22 and the pExpAD502 plasmids were used as

negative controls. All Y2H procedures were performed following the manufacturer's instructions. Quantitation of β -galactosidase activity was performed in a 96-well format as previously described (Pruneda-Paz et al., 2009).

In Vitro Pull-Downs

For *in vitro* pull-down assays, additional constructs were made. The pENTR/D-TOPO plasmids (Invitrogen) containing the sequences encoding GI, PIF1, and PIF5 have been previously described (Pruneda-Paz et al., 2014; Sawa and Kay, 2011). Full-length *PIF3*, *PIF4*, and *GFP* sequences were amplified by PCR (primers used are listed in Table S2) and cloned into the pENTR/D-TOPO vector (Invitrogen). Partial *PIF3* sequences were amplified by PCR (primers used are listed in Table S2) and cloned into the pDONRZeo vector (Invitrogen) by Gateway BP recombination reaction (Invitrogen). To express proteins in the cell-free system, all inserts were transferred by Gateway LR recombination reaction (Invitrogen) into Gateway compatible modified pTnT vectors (Promega) (Nito et al., 2013), which were kindly provided by Dr. Joanne Chory (The SALK Institute, La Jolla, CA). The vectors contained an N-terminal HA or Flag tag as specified in each case. Proteins were co-expressed using TnT SP6 High-Yield Wheat Germ Protein Expression System (Promega) as per manufacturer's instructions. Five percent of the reactions (2.5 μ l) were used to verify expression of the proteins (input) and the remaining extract was immunoprecipitated as earlier described (Pedmale et al., 2016) using anti-HA 3F10 antibody (Roche).

Transient Expression in *N. benthamiana*

Agrobacterium tumefaciens strain GV3101 was used in all instances, except for the constructs derived from the pGreenII 0800-LUC vector, which were transformed into the *A. tumefaciens* strain C58 containing the pSoup helper plasmid. *A. tumefaciens* cells containing the respective constructs and the p19 silencing suppressor were grown overnight at 28°C in liquid LB medium supplemented with the appropriate antibiotics. Cultures were pelleted, resuspended in 10 mM MES-KOH pH 5.6, 10 mM MgCl₂, 150 μ M acetosyringone to a final OD₆₀₀ of 0.5, and incubated for 2 h at room temperature. The suspensions were then mixed and infiltrated in *N. benthamiana* leaves at a final OD₆₀₀=0.1 each, except for p19 which was infiltrated at a final OD₆₀₀=0.05. Samples were harvested 2 days post-inoculation (dpi) for transactivation assays and 3 dpi for co-localization and protein stability analyses. In the case of MG-132 treatments, plants were infiltrated with 25 μ M MG-132 8 h before harvesting.

Protein Immunoprecipitation

Approximately 1 g of 10-day-old *Arabidopsis* seedlings grown in SDs were harvested at ZT 8 and frozen in liquid nitrogen. For time-course experiments, 0.5 g were used. Immunoprecipitations were performed as earlier described (Nusinow et al., 2011) with the following modifications. Samples were ground with mortar and pestle in liquid nitrogen and resuspended in 2 ml of modified SII buffer (100 mM NaPhosphate, pH 8.0, 150 mM NaCl, 5 mM EDTA, 5 mM EGTA, 0.1% Triton X-100, 2 mM PMSF, 1x protease inhibitor cocktail (Roche), 1x Phosphatase Inhibitors 2&3 (Sigma) and 50 μ M MG-132 (Peptides International)). Extracts were transferred to a dounce tissue grinder and homogenized before being clarified twice by centrifugation at 4°C. Total protein concentration was quantified by DC Protein Assay (Bio-Rad, Hercules, California, United States of America), and normalized to 1.875 mg/ml. Three percent of the extracts was used to verify proteins levels (input). For immunoprecipitation, extracts were incubated with anti-Flag M2 antibody (Sigma) for 2 h with gentle rotation at 4°C. Subsequently, 25 μ l of magnetic protein G Dynabeads (Invitrogen) pre-washed with IP buffer were added to the samples and incubated for 2 h with gentle rotation at 4°C. The samples were finally washed 3x with modified SII buffer and the precipitated protein was eluted by heating beads at 95°C for 5 min in 40 μ l of 2x SDS-PAGE loading buffer. 10 and 30 μ l of the eluate were separately analyzed by western blot to detect the immunoprecipitated and co-immunoprecipitated proteins, respectively.

Protein Stability and Accumulation Analyses

To determine protein levels in transient experiments in *N. benthamiana*, samples were homogenized with 3 volumes of 2x SDS-PAGE loading buffer and boiled at 95°C for 5 min. Samples were then clarified by centrifugation at room temperature and analyzed by western blot. For normalization, GFP-HA or p19-HA were used as internal loading controls. Alkaline phosphatase treatments were performed as earlier described (Ni et al., 2013) with the following modifications. Proteins were extracted with 2x SDS-PAGE loading buffer, boiled at 95°C for 5 min, and clarified by centrifugation. Extracted proteins were diluted 10-fold into reaction buffer (50 mM Tris-HCl pH 8.8, 100 mM NaCl, 50 mM MgCl₂, 1% Triton X-100, 1x protease inhibitor cocktail (Roche)) and incubated in the presence or absence of CIP (400 U/ml) for 2 h at 37°C. Samples were then analyzed by western blot.

For the analysis of PIF3-ECFP-HA and PIF5-HA protein levels in *Arabidopsis* seedlings, samples were homogenized with 100 μ l of extraction buffer (50 mM Tris-HCl pH 7.6, 150 mM NaCl, 5 mM MgCl₂, 0.1% NP-40, 10% glycerol, 2 mM PMSF, 1x protease inhibitor cocktail (Roche), 1x Phosphatase Inhibitors 2&3 (Sigma) and 50 μ M MG-132 (Peptides International)) and clarified twice by centrifugation at 4°C. Total protein concentration was quantified by DC Protein Assay (Bio-Rad) and 40 μ g of each sample was subsequently analyzed by western blot. ACTIN levels in the samples were used for normalization.

Western Blot Detection and Quantitation

Protein extracts in SDS-PAGE loading buffer were boiled at 95°C for 5 min and separated in 4–15 % SDS-PAGE gels. Proteins were then transferred to nitrocellulose membranes (BioRad), which were then stained with Ponceau S to assess transfer and loading. Finally, membranes were immunodetected with either Horse Radish Peroxidase (HRP)-conjugated 3F10 anti-HA (1:2000, Roche)

or HRP-conjugated Flag M2 (1:2000, Sigma) antibody. The ACTIN loading control was detected using anti-ACTIN C4 mouse antibody (1:500, Millipore) followed by HRP-conjugated anti-mouse secondary antibody (1:3000, BioRad). *Arabidopsis* PIF3-ECFP-HA and PIF5-HA blots were cut at the 50 kDa mark and the upper and lower parts were detected with anti-HA and anti-ACTIN antibodies, respectively. Chemiluminescence was detected with the Supersignal West Pico, Dura, and Femto substrates (Pierce) and imaged with a UVP ChemiDoc imaging system (UVP, LLC) (in which case the VisionWorksLS (UVP, LLC) software was used to quantify protein levels) or captured by exposure of X-ray films (GE Healthcare), which were scanned and protein levels quantified using NIH ImageJ software (Schneider et al., 2012) (<https://imagej.nih.gov/ij/>).

Electrophoretic Mobility Shift Assay (EMSA)

DNA probes for gel-shift assays (Martínez-García et al., 2000) were generated by annealing the oligonucleotides listed in Table S2 (N629+N630 for the labeled probe, and N602+N603 for the unlabeled one). Proteins were produced by *in vitro* transcription and translation (Martínez-García et al., 2000) using the TnT SP6 High-Yield Wheat Germ Protein Expression System (Promega) as per manufacturer's instructions. Binding reactions were prepared by mixing the indicated amounts of protein solutions in binding buffer (10 mM Tris-HCl pH 7.5, 50 mM KCl, 1 mM DTT, 5 mM MgCl₂, 2.5% glycerol, 0.05% NP-40) containing 1 µg poly (dI-dC) and 1 µl of the end-labeled probe (100 nM) in a final volume of 20 µl. For competitions, the indicated amounts of unlabeled probe or protein extract were added to the reaction. After incubation for 20 minutes at room temperature, complexes were resolved by 6% PAGE in 0.5x TBE buffer at room temperature. Gels were scanned with a Typhoon 9410 imager (GE Healthcare).

RNA Extraction and qRT-PCR

Total RNA was isolated with the GeneJET plant RNA purification mini kit (Thermo Scientific). For cDNA synthesis, 1 µg of total RNA was digested with DNase I (Roche) and reverse-transcribed using the iScript cDNA synthesis kit (Bio-Rad). Synthesized cDNA was amplified by real-time quantitative PCR (qPCR) with Maxima SYBR Green qPCR Master Mix (Thermo Scientific) using the CFX-384 Real Time System (BioRad). *PROTEIN PHOSPHATASE 2A (PP2A)* (AT1G13320) was used as the normalization control. Primer sequences are listed in Table S2.

Transactivation Assays

Reporter and effector constructs were co-infiltrated in *N. benthamiana* leaves and 2 dpi transient activation assays were performed as earlier described (Nieto et al., 2015).

Chromatin Immunoprecipitation (ChIP)

Approximately 4 g of fresh weight whole seedlings were harvested at the indicated ZTs and cross-linked with 1% formaldehyde under vacuum for 10 min. Cross-linking was quenched by adding glycine to a final concentration of 125 mM, pH 8.0 under vacuum for 5 min. Seedlings were then washed three times in ddH₂O and rapidly frozen in liquid nitrogen. Samples were ground with mortar and pestle in liquid nitrogen and processed as described (Sawa et al., 2007) with the following modifications: lysed nuclei were diluted 4-fold with ChIP dilution buffer (16.7 mM Tris-HCl pH 8, 167 mM NaCl, 1.1% Triton X-100, 1.2 mM EDTA, 1 mM PMSF, 5 mM Benzamide, 1x protease inhibitor cocktail (Roche)) and sonicated using a Bioruptor (Diagenode) (0.5 s on/off cycles for 15 min total at maximum power) to shear DNA to an average size of 500-1000 bp. After sonication, samples were clarified twice by centrifugation at 4°C and the chromatin solution was diluted 2-fold with ChIP dilution buffer and pre-cleared by adding 50 µl of magnetic protein G Dynabeads (Invitrogen) pre-washed with ChIP dilution buffer for 1 h with gentle rotation at 4°C. At this point, 100 µl were saved as input. Immunoprecipitation was then performed overnight with 4 µg of anti-HA 3F10 (Roche) or anti-Flag M2 (Sigma) antibody at 4°C with gentle rotation. Mock samples were processed without antibody. Subsequently, 50 µl of magnetic protein G Dynabeads (Invitrogen) pre-washed with ChIP dilution buffer were added and samples were incubated for 2 h with gentle rotation at 4°C. Low and high salt washes were followed by LiCl buffer (250 mM LiCl, 1% Sodium Deoxycholate, 1% NP-40, 1 mM EDTA) and TE buffer (10 mM Tris-HCl pH 8, 1 mM EDTA) washes. Immunocomplexes were eluted twice from the beads using a modified elution buffer (100 mM NaHCO₃, 10 mM EDTA, 1% SDS) and incubation at 65°C. Crosslinking in the immunoprecipitated and input samples was reversed by adding NaCl to a final concentration of 200 mM and incubating at 65°C overnight, which was followed by a treatment with Proteinase K (Invitrogen) to degrade all proteins. DNA was purified by phenol-chloroform extraction and ethanol precipitation. DNA was resuspended in 75 µl of TE and 1 µl aliquots were used for qPCR. For the input samples, 1 µl of a 100-fold dilution was used for qPCR. qPCR reactions were performed in triplicate with Maxima SYBR Green qPCR Master Mix (Thermo Scientific) using the Stratagene Mx3005P qPCR system (Agilent Technologies). Primers used were as in previous studies (Johnson et al., 2014; Nieto et al., 2015; Soy et al., 2016) and are listed in Table S2. Enrichment in each sample was calculated relative to the input and expressed in percentage.

ChIP-seq

ChIP-seq libraries were prepared with the NuGEN Ovation Ultralow System as per manufacturer's instructions. For quantitative ChIP-seq, 0.02 ng of sonicated unmethylated lambda DNA (Promega) were added to each sample prior to library preparation.

Physiological Measurements

To analyze hypocotyl length, evenly spaced seedlings were grown on plates under the indicated light conditions and photoperiod. At the specified time, seedlings were scanned and images were analyzed using NIH ImageJ software (<https://imagej.nih.gov/ij/>).

For growth rate analyses, plants were grown and analyzed essentially as described (Nozue et al., 2007) except that the growth media was 1/2X MSMO (Sigma), 0.8% phytagar (Sigma) and images were captured with a PixelINK PL-A781 camera driven by LabView (National Instruments). Plants were grown under short day cycles (8h light, 16 h dark) at 22°C in a Conviron E7 plant growth chamber. 72 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetically active radiation was provided by cool white fluorescent bulbs.

Clock Activity Measurements

For circadian rhythm measurements, seedlings were grown on 96-well plates and entrained in SD conditions. After 7 days of entrainment, 500 μM luciferin (Biosynth AG) was added to the wells and the plates were transferred to constant light (80 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Bioluminescence was recorded for 5 days. Diel monitoring of *pCCA1::LUC* expression was conducted similarly, except that seedlings were kept in SDs while recording over the course of 2 days. In order to analyze the light-induced activation of *pCCA1*, etiolated seedlings were grown on 96-well plates for 3 days in constant darkness after stratification and germination induction. Bioluminescence started to be recorded immediately after transfer to constant white light. Light-induced phase shifts of *pCCA1::LUC* expression were quantified as earlier described (Viczián et al., 2005). Briefly, seedlings were grown under SDs for 5 days, and then kept in continuous darkness for 24 h. After 24 h in darkness, plates were irradiated with white light at the indicated ZTs for 1 h and then transferred to the luminometer to monitor luminescence. Phase shifts were calculated by comparing the phase values of the pulsed plants with those of non-pulsed controls. In all cases, bioluminescence was recorded every 1 h over the indicated period of time with a Tecan Infinite M200 plate reader (with the exception of Figures 4E and S2G - PIF1 and PIF4 panels, for which a Centro LB960 Microplate Luminometer (Berthold Technologies) was used due to logistic reasons) and data were analyzed by fast Fourier transformed non-linear least squares (FFT-NLLS) using BioDare2 (Moore et al., 2014; Zielinski et al., 2014) (<https://www.biodare2.ed.ac.uk>). For phase shift calculations, the LS Periodogram algorithm was applied (Zielinski et al., 2014).

QUANTIFICATION AND STATISTICAL ANALYSIS

Mean Comparisons

In order to test the statistical significance of the difference between mean values relative to wildtype controls, Student's *t* test and Tukey's multiple comparisons test were applied, as specified in each experiment. Sample size and number of replicates (where applicable) for each experiment are indicated in the respective figure legend. Information about types of tissue can be found in the respective figure legend and in the STAR Methods Details section. Strategies for randomization and/or stratification and sample size estimation were not applied to these experiments. No statistical calculation was used to estimate the sample size.

ChIP-seq Data Analysis

Raw sequencing reads from the sequencer were demultiplexed and quality controlled with FastQC (v0.11.3). Reads with high quality were then aligned to the Arabidopsis reference genome (TAIR10) with Bowtie (v1.0.0) (Langmead et al., 2009), allowing only uniquely mapping reads with fewer than two mismatches, and duplicated reads were combined into one read. For quantitative ChIP-seq, reads were also mapped to lambda DNA with Bowtie (v1.0.0) (Langmead et al., 2009) with the same parameters as before. For both Flag and HA ChIP-seqs, peaks were called with MACS2 (v 2.1.1.) (Zhang et al., 2008) using Flag and HA ChIP-seqs in Col-0 as controls, respectively. The set of peaks identified from both the GI and PIF3 ChIP-seqs can be found in the Table S1 file. In all cases, ChIP-seq signal was measured in reads per kilobase million (RPKM), and normalized to the signal in the control Col-0 IP. For quantitative ChIP-seq, the signal was additionally normalized to the lambda DNA spike-in control. ChIP-seq metaplots were plotted using NGSplot (v 2.41.4) (Shen et al., 2014). *De novo* predominant motifs analysis was conducted using HOMER (Heinz et al., 2010) over a 200 bp region around the ChIP-seq peak summits.

Genome-Wide Gene Expression Analyses

All datasets used for genome-wide gene expression analysis are compiled in the Table S1.

The statistical significance of the overlap between gene lists was calculated using Fisher's exact test.

Identification of over-represented known plant transcription factor binding motifs at 1000 bp promoter regions of DEGs in *gi-2* seedlings (Kim et al., 2012) was performed with HOMER (Heinz et al., 2010).

The web-based tool Phaser (<http://phaser.mocklerlab.org/>) (Mockler et al., 2007) was used to determine the phases of peak gene expression and their over-representation.

For the identification of over-represented Gene Ontology (GO) categories in gene lists, the web-based tool PANTHER (<http://www.pantherdb.org/>) (Thomas et al., 2003) was used. For the significantly enriched categories (*p* value < 0.05) in each subset, enrichment scores were calculated as $-\log_{10}(p)$, where *p* is the *p* value from the enrichment analysis.

Growth Rate Calculations

Change in growth was calculated for each 30-min time interval and local polynomial regression fitting (loess) smoothing with smoothing parameter = 0.1 was used in R (R Development Core Team, 2016) and results were plotted with the ggplot2 plotting package (Wickham, 2009). Analysis scripts are available at <https://github.com/MaloofLab/Nohales-2019>.

DATA AND SOFTWARE AVAILABILITY

The accession number for the high-throughput sequencing data reported in this paper is GEO: GSE129865.

Developmental Cell, Volume 49

Supplemental Information

Multi-level Modulation of Light

Signaling by GIGANTEA Regulates

Both the Output and Pace of the Circadian Clock

Maria A. Nohales, Wanlu Liu, Tomas Duffy, Kazunari Nozue, Mariko Sawa, Jose L. Pruneda-Paz, Julin N. Maloof, Steven E. Jacobsen, and Steve A. Kay




Figure S1. GI shares targets with light signaling components and interacts with PIF proteins, Related to Figure 1

(A) Overlap between DEGs in *gi-2*, a comprehensive set of PIF-regulated genes, and DEGs in *cca1-1;lhy-11* mutants (GI-PIF intersection $p < 2.2 \times 10^{-16}$; GI-CCA1/LHY intersection $p < 2.2 \times 10^{-16}$).

(B and C) Phase enrichment heatmaps depicting the p value of the phase of peak expression enrichment of genes differentially expressed in *gi-2* only (GI), regulated by PIFs only (PIFs), and potentially regulated by both GI and PIFs (GI-PIFs) (B) or DEGs in *gi-2* only (GI), *cca1-1;lhy-11* only (CCA1;LHY), and in both *gi-2* and *cca1-1;lhy-11* (GI/CCA1;LHY) (C) under 12h light and 12h dark conditions (* $p < 0.01$). Day period is marked in yellow and night period in gray.

(D) Heat map showing the GO term enrichment scores of genes differentially expressed in *gi-2* only (GI), potentially regulated by both GI and PIFs (GI-PIFs), and differentially expressed in both *gi-2* and *cca1-1;lhy-11* (GI-CCA1/LHY).

(E) Hypocotyl length measurements from wildtype (Col-0) and *gi-2* seedlings grown for 6 days under different light conditions (in gray, mean $\pm$ SEM, $n = 20-36$; *** $p < 0.001$ Student's *t*-test). D, darkness; cR, constant red light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$); cB, constant blue light ($1 \mu\text{mol m}^{-2}\text{s}^{-1}$); SD, short day photoperiod (8h light, 16h dark).

(F, G) Yeast two-hybrid (Y2H) assays showing interaction of GI and PIF proteins. Bait and prey constructs were co-transformed into yeast cells. SD-WL, minimal medium lacking Trp and Leu; SD-WLH, selective medium lacking Trp, Leu and His, which was supplemented with 50 mM 3AT; X-gal, qualitative β -galactosidase activity results obtained from the X-gal assay. (G) Quantitation of β -galactosidase activity (Miller units) for every pair of bait and prey proteins indicated ($n = 4$). Values represent means $\pm$ SEM. Statistically significant differences between mean values by Student's *t*-test relative to the pExpAD502 control vector are shown (** $p < 0.01$, *** $p < 0.001$).

(H) *In vitro* pull-down assays performed to map the interaction domains between GI and PIF3. Proteins were expressed in a TnT *in vitro* expression system and immunoprecipitated with anti-HA antibody. The recovered fractions were analyzed by Western blot using anti-Flag and anti-HA antibodies. A scheme of the deleted protein versions is shown on the upper panel.




Figure S2. GI modulates PIF stability and activity, Related to Figure 2

(A and B) Relative expression of *HFR1* and *PRE5* (A) and *PIF1*, *PIF3*, *PIF4*, and *PIF5* (B) in wildtype (Col-0), *gi-2*, and Glox seedlings grown for 10 days in SDs (mean $\pm$ SEM of 3 biological replicates). White and gray shadings represent day and night, respectively.

(C and D) HA-PIF3 (C) and HA-PIF5 (D) accumulation in *N. benthamiana* leaves in the presence or absence of Flag-GI. Protein levels were normalized against HA-GFP levels. Western blot quantitation is shown on the respective lower panels. Values represent mean $\pm$ SEM (n=3).

(E) Western blot analysis of the treatment with calf intestinal alkaline phosphatase (CIP) of PIF5-HA expressed in *N. benthamiana* leaves in the presence and absence of GI. Western blot quantitation is shown on the lower panel. Values represent mean $\pm$ SEM (n=2).

(F) Relative expression of the *PIF5-HA* transgene in wildtype (Col-0) and Glox plants at ZT8 and 16 in SDs (mean $\pm$ SEM of 3 biological replicates).

(G) Transactivation assays in *N. benthamiana* leaves. Different effectors were co-expressed with the *pPIL1::LUC* reporter construct. Luminescence was measured 2 days post-infiltration in SD conditions. Results show mean $\pm$ SEM (n=12).

(H, J and K) EMSA studies using a *PIL1* promoter fragment containing two G-boxes as probe. Flag-PIF3 (H), HA-GI (H,J,K), and Flag-PIF5 (K) were expressed in an *in vitro* transcription and translation system. The Cy5 labeled DNA probe was incubated with increasing amounts of the proteins as indicated (1, 2, 4 μ l) or with the maximal amount of protein and increasing quantities of unlabeled probe (50, 100, 200 fold excess) (competitor, C). E, control incubation with 4 μ l extract without expressed protein. 0, incubation without extract. M, molecular weight marker. (H and K) The binding of Flag-PIF3 and Flag-PIF5 to the probe was additionally analyzed in the presence of increasing quantities of HA-GI (1, 2, 4 μ l).

(I) Western blot analysis of Flag-PIF3 stability in the presence of increasing quantities of HA-GI (1, 2, 4 μ l) during the binding reactions.

(A-F) n.s. not significant, *p<0.05, **p<0.01, ***p<0.001 Student's *t*-test.




Figure S3. GI modulates PIF stability and activity, Related to Figure 2

(A) HA-PIF3 accumulation in *N. benthamiana* leaves treated with 25 μ M MG-132 or in the presence of Flag-GI. Protein levels were normalized against HA-GFP levels. Western blot quantitation is shown on the lower panel. Values represent mean $\pm$ SEM (n=3).

(B) PIF3-ECFP-HA protein accumulation at ZT8 and 16 in the indicated backgrounds under SD conditions. Protein levels were normalized against ACTIN levels. The quantitation is shown on the lower panel (mean $\pm$ SEM of 3 biological replicates).

(C) Relative expression of the *PIF3-ECFP-HA* transgene in wildtype (Col-0) and Glox plants at ZT8 and 16 in SDs (mean $\pm$ SEM of 3 biological replicates).

(D and E) PIF3-ECFP-HA (D) and PIF5-HA (E) accumulation in Col-0 and Glox plants in etiolated seedlings grown for 3 days with (PAC) or without (D) 1 μ M paclobutrazol. Protein levels normalized against ACTIN levels. The quantitations are shown on the respective lower panels. Values represent mean $\pm$ SEM (n=3).

(F) Physiological characterization of PIF3-ECFP-HA overexpression lines. Left panel, hypocotyl length measurements (n=14-17) from wildtype (Col-0) and PIF3-ECFP-HA seedlings grown for 7 days in constant white light (WL), SDs (SD) or for 3 days in constant red light (Rc, 10 μ mol m²s⁻¹). Right panel, relative expression of *PIL1* in wildtype (Col-0) and PIF3-ECFP-HA seedlings at the indicated ZTs in SDs (mean $\pm$ SEM of 3 biological replicates).

(G) Representative western blots showing the accumulation of PIF3-ECFP-HA across a SD photo-cycle in the wildtype background (Col-0) and in Glox plants. Expression of PIF3-ECFP-HA in these lines is driven by the *PIF3* endogenous promoter. Protein levels were determined with anti-HA antibody and ACTIN levels were used as loading control. Western blot quantitation is shown on the right panel (mean $\pm$ SEM, n=3 biological replicates).

(H and I) PIF3 (H left panel and I) and PIF5 (H right panel) ChIP assays of 10-day-old seedlings grown in SD conditions and harvested at ZT8 in the indicated lines. The enrichment of the specified regions in the immunoprecipitated samples was quantified by qPCR. Values represent mean $\pm$ SEM (n=2-4).

(J) Relative expression of *PIL1* in 10-day-old seedlings of the indicated backgrounds and ZTs in SDs (mean $\pm$ SEM of 3 biological replicates). White and gray shadings represent day and night, respectively.

(K) Relative expression of *PIL1* and *HFR1* was additionally analyzed in 3-day-old seedlings of the indicated backgrounds in SDs (mean $\pm$ SEM of 3 biological replicates).

(L) Relative expression of *PSY* in 10-day-old seedlings of the indicated backgrounds and ZTs in SDs (mean $\pm$ SEM of 3 biological replicates). White and gray shadings represent day and night, respectively.

(A-L) n.s. not significant, *p<0.05, **p<0.01, ***p<0.001 Student's *t*-test.




Figure S4. GI and PIFs occupy the same genomic targets in a phase dependent pattern, Related to Figure 3

(A) Protein accumulation pattern of GI-YPET-Flag (driven by an endogenous promoter fragment), PIF3-ECFP-HA (driven by an endogenous promoter fragment), and PIF5-HA (overexpressed) across SD photo-cycles. Protein levels were quantified relative to ACTIN levels.

(B and C) Time-course *in vivo* co-immunoprecipitation assays in *Arabidopsis* transgenic seedlings expressing GI-YPET-Flag and PIF3-ECFP-HA (driven by endogenous promoter fragments) (B, left panel), and GI-YPET-Flag and PIF5-HA (overexpressed) (B, right panel, and C) tagged protein versions.

(D) Overlap between peaks called in the different GI ChIP-seq biological replicates. Peaks identified in all three replicates were considered high confidence GI target sites (n=7365).

(E) Genomic annotation of a random permuted set of similar sample size to GI ChIP-seq peaks. Control to Figure 3D.

(F) Genomic annotation of PIF3 (n=7962) ChIP-seq peaks (left panel) compared to a random permuted set of similar sample size (right panel). The midpoint of the peaks was used for this analysis.

(G) Over-represented *cis* elements around the summit of PIF3 ChIP-seq peaks (± 100 bp flanking region) in SD conditions.

(H) Heat map showing the GO term enrichment scores of genomic targets shared by GI and PIF3.

(I) Phase enrichment graph depicting the enrichment (count/expected) in the phase of peak expression of genomic targets shared by GI and PIF3 in SDs (* $p < 0.01$). Day period is marked in white and night period in gray.

(J) Visualization of PIF3 and GI ChIP-seq data in the genomic region encompassing the *HFR1* locus.

(K) Boxplot of the signal from GI ChIP-seq peaks ranked by increasing signal and divided in 10 groups (deciles 1 to 10).

(L) Growth rate measurements (mm/h) of wildtype (Col-0), *gi-2*, *pif3-1*, and *gi-2;pif3-1* seedlings grown in SDs. Values represent mean $\pm$ SEM (n=18-36). Time is expressed in hours after stratification. White and gray shadings represent day and night, respectively. The right panel shows the overlap of the growth rate measurement traces of *gi-2* and *gi-2;pif3-1* seedlings.




Figure S5. GI modulation of light signaling affects circadian rhythms, Related to Figure 4

(A) Transactivation assays in *N. benthamiana* leaves. Different effectors were co-expressed with the *pCCA1::LUC* reporter construct. Luminescence was measured 2 days post-infiltration in SD conditions. Results show mean $\pm$ SEM (n=8).

(B) Relative expression of *CCA1* in the indicated backgrounds and ZTs in SDs (mean $\pm$ SEM of 3 biological replicates).

(C) Bioluminescence analysis of *pCCA1::LUC* in seedlings in SD conditions. Values represent mean $\pm$ SEM (n=12). White and gray bars represent day and night, respectively.

(D and F) Bioluminescence analysis of *pCCA1::LUC* in seedlings of the indicated backgrounds in constant light (LL). Values represent mean $\pm$ SEM (n=24). Plants were entrained in SDs for 7 days.

(E and G) Left panels, period length estimations versus RAE of *pCCA1::LUC* in D and F respectively as analyzed by FTT-NLLS. Right panels, period length estimations of *pCCA1::LUC* in D and F respectively as analyzed by FTT-NLLS. Statistically significant differences between mean values by Student's *t*-test relative to the Col-0 control are shown (**p<0.01, ***p<0.001).

(H and J) Data shown in Figure 4E and G respectively is re-plotted on a different scale for the y-axis.

(I and K) Period length estimations versus RAE of *pCCA1::LUC* data shown in Figure 4E and G respectively as analyzed by FTT-NLLS.

(L) Bioluminescence analysis of *pCCA1::LUC* in 3-day-old etiolated seedlings transferred to light (mean $\pm$ SEM, n=12).

(M) Light-induced phase shifts of *pCCA1::LUC* expression plotted against the ZT at which the light pulse was given. Positive values represent phase advances and negative values, phase delays. Values represent mean $\pm$ SEM (n=6).