

# Epigenetic maintenance of the vernalized state in *Arabidopsis thaliana* requires LIKE HETEROCHROMATIN PROTEIN 1

Sibum Sung<sup>1,5</sup>, Yuehui He<sup>1,4,5</sup>, Tifani W Eshoo<sup>1</sup>, Yosuke Tamada<sup>1</sup>, Lianna Johnson<sup>2</sup>, Kenji Nakahigashi<sup>3,4</sup>, Koji Goto<sup>3,4</sup>, Steve E Jacobsen<sup>2</sup> & Richard M Amasino<sup>1</sup>

**Vernalization is the process by which sensing a prolonged exposure to winter cold leads to competence to flower in the spring. In winter annual *Arabidopsis thaliana* accessions, flowering is suppressed in the fall by expression of the potent floral repressor FLOWERING LOCUS C (FLC)<sup>1</sup>. Vernalization promotes flowering via epigenetic repression of FLC<sup>2</sup>. Repression is accompanied by a series of histone modifications of FLC chromatin that include dimethylation of histone H3 at Lys9 (H3K9) and Lys27 (H3K27)<sup>3,4</sup>. Here, we report that *A. thaliana* LIKE HETEROCHROMATIN PROTEIN 1 (LHP1) is necessary to maintain the epigenetically repressed state of FLC upon return to warm conditions typical of spring. LHP1 is enriched at FLC chromatin after prolonged exposure to cold, and LHP1 activity is needed to maintain the increased levels of H3K9 dimethylation at FLC chromatin that are characteristic of the vernalized state.**

**ngp** In many plant species, the timing of flowering is affected by seasonal cues. Plants that flower early in the spring often have a vernalization requirement; such plants require prolonged exposure to winter cold in order to achieve competence to flower. That vernalization typically requires prolonged cold permits plants to discriminate between transient warm periods in the fall and the actual arrival of spring<sup>2</sup>.

In *A. thaliana*, vernalization promotes flowering primarily by repressing the flowering repressor FLOWERING LOCUS C (FLC)<sup>5,6</sup>. Vernalization-mediated FLC repression is epigenetic in the sense that it is maintained throughout subsequent mitotic cell divisions in the shoot apical meristem during spring growth. Expression of FLC resumes in the next generation, thus perpetuating the winter-annual growth habit<sup>1</sup>.


Vernalization-mediated FLC repression is accompanied by deacetylation of H3K9 and H3K14 and methylation of H3K9 and H3K27 (refs. 3,4), which are likely to result in formation of a mitotically stable

heterochromatin-like state at FLC. Maintenance of FLC repression requires VERNALIZATION2 (VRN2), the *A. thaliana* homolog of Suppressor of zeste 12 (Su(z)12)<sup>7</sup>. In animals, Su(z)12 is a component of Polycomb repression complex 2 (PRC2), a complex involved in maintaining Polycomb group (PcG) repression of target genes through methylation of H3K27 (ref. 8). VRN2 is likely to have a role similar to Su(z)12 in FLC repression, because in *vrn2* mutants the level of H3K27 methylation of FLC chromatin does not increase during cold exposure, and FLC repression is not maintained in subsequent cell division cycles<sup>3,4</sup>. In addition, *A. thaliana* has other PRC2 components; for example, there are three homologs of the H3K27 methyltransferase Enhancer of zeste (E(z))<sup>9</sup>. Thus, both animals and plants seem to use PRC2-type epigenetic repression involving H3K27 methylation.

In animals, maintenance of PRC2-mediated repression requires Polycomb repression complex 1 (PRC1)<sup>8</sup>. When H3K27 is methylated, PRC1 binds to histone H3 via the chromodomain of one of its components, Polycomb protein (Pc)<sup>8</sup>. Notably, no homologs of PRC1 components have been identified in plants<sup>9</sup>. Therefore, plants are likely to have a system different from that of animals for the maintenance of PcG-type silencing. Indeed, in animals, methylation of H3K27 seems to be a hallmark of genes that are stably repressed by PcG-mediated gene silencing<sup>8</sup>, but the stable repression of FLC by vernalization requires methylation of not only H3K27 but also H3K9 (refs. 3,4).

One component of the system that might have a PRC1-like role in plants is the homolog of HETEROCHROMATIN PROTEIN 1 (HP1), LIKE HETEROCHROMATIN PROTEIN 1 (LHP1; also called TERMINAL FLOWER 2)<sup>10,11</sup>. In animals and yeast, HP1 preferentially localizes to, and is involved in maintenance of, constitutive heterochromatin, such as pericentric regions<sup>12</sup>, although there are examples of a role for HP1 in the regulation of euchromatic genes in *Drosophila melanogaster* and mammals<sup>13,14</sup>. HP1 binds to chromatin by interacting with methylated H3K9 (ref. 12), a modification associated with maintenance of the vernalized state<sup>3,4</sup>. In plants, LHP1 localizes

<sup>1</sup>Department of Biochemistry, University of Wisconsin at Madison, Madison, Wisconsin, USA. <sup>2</sup>Howard Hughes Medical Institute, Department of Molecular, Cell, and Developmental Biology, and Molecular Biology Institute, University of California, Los Angeles, California 90095, USA. <sup>3</sup>Research Institute for Biological Sciences, Okayama, 716-1241 Japan. <sup>4</sup>Present addresses: Department of Biological Sciences, Faculty of Science, National University of Singapore, Science Drive 4, Singapore 117543 (Y.H.); Institute for Advanced Biosciences, Yamagata 997-0017, Japan (K.N. and K.G.). <sup>5</sup>These authors contributed equally to this work. Correspondence should be addressed to R.M.A. (amasino@biochem.wisc.edu).



**Figure 1** The vernalization response is impaired in *lhp1* mutants. **(a)** Flowering behavior of *lhp1* mutants with and without vernalization in long days and short days. Flowering was measured as the number of rosette leaves produced from primary meristems prior to flowering. Arrows at tops of bars indicate that plants had not flowered after producing more than 100 primary rosette leaves. **(b)** *lhp1* lesion blocks the vernalization response. *FRI*-containing winter annual lines without (upper left) and with (upper right) vernalization in short days. *lhp1-4l/FRI-Col* lines without (lower left) and with (lower right) vernalization in short days. **(c)** mRNA levels of *FLC* and *FT* in *lhp1-4l/FRI-Col*. *FT* levels are elevated in *lhp1-4l/FRI-Col*, causing early flowering in long days despite higher levels of *FLC*. However, *FT* is only transiently expressed when *FLC* is transiently repressed during vernalization in short days. **(d)** Repression of *FLC* is not maintained in *lhp1-4l/FRI-Col*.

days. Specifically, the decreased CO activity in short days may shift the balance of promotive and repressive activities such that *FLC* is able to repress *FT* in short days in the absence of

primarily to euchromatic regions (Supplementary Fig. 1 online)<sup>15–17</sup>, similar to the distribution of PRC1 in animals<sup>8</sup>. Furthermore, LHP1 does not seem to be required for repression of loci that are silenced by DNA methylation, including pericentric regions, but it may have a direct role in the stable repression of euchromatic genes<sup>15,18</sup>.


To evaluate whether *LHP1* has a role in vernalization, we introduced the *lhp1* mutation into the vernalization-responsive genetic background, *FRI-Col* (flowering of *FRI-Col* plants is extremely delayed in inductive photoperiods, unless the plants are vernalized<sup>19</sup> (Fig. 1a)). In long days without exposure to cold, however, the *lhp1* lesion in *FRI-Col* (*lhp1/FRI-Col*) causes rapid flowering (Fig. 1a); therefore, growth in long days precludes assessing the vernalization response. In contrast, without vernalization, *FRI-Col* plants with or without the *lhp1* lesion often fail to flower in noninductive photoperiods (that is, short days; Fig. 1a,b). Thus, growth in short days provides a means to assess the role of *LHP1* in vernalization. In short days, cold treatment strongly promotes flowering of the *FRI-Col* line but has no effect on the flowering of *lhp1/FRI-Col* (Fig. 1a,b), demonstrating that *LHP1* is required for the vernalization response.

To determine which phase of *FLC* silencing requires *LHP1*, we monitored *FLC* levels during and after cold exposure. *FLC* mRNA levels declined during cold exposure in *lhp1/FRI-Col*, similar to wild-type plants, but after we returned *lhp1/FRI-Col* plants to warm conditions, *FLC* mRNA increased to prevernalization levels (Fig. 1c,d)<sup>20</sup>. Thus, *LHP1* is required for the maintenance of *FLC* repression.


The ability of the *lhp1* mutation to suppress the *FLC*-mediated delay of flowering in long days but not in short days (Fig. 1a) is associated with differential expression of the floral promoter *FT* in the different photoperiods (Fig. 1c). In long days, *FT* expression is elevated in *lhp1/FRI-Col* plants despite high levels of *FLC*; in contrast, in short days, *FT* levels are low. In the absence of elevated *FLC* expression, *lhp1* lesions result in early flowering and elevated *FT* expression in short days<sup>11,15</sup>. Thus, *FLC* can repress *FT* in short days but not in long days in *lhp1* mutants. This photoperiod-dependent *FLC* repression of *FT* is likely to result from the differential activity of *CONSTANS* (CO), a promoter of *FT* expression<sup>21</sup>, in long and short

*LHP1*. Notably, in *lhp1/FRI-Col*, *FT* is only expressed while *FLC* is transiently repressed (Fig. 1c), indicating that the resumption of *FLC* expression may prevent the positive autoregulation of *FT* that was recently reported to occur in a rapid-cycling genetic background that does not express *FLC* to levels that repress flowering<sup>22</sup>.

To test whether the association of *LHP1* with *FLC* chromatin is enriched by vernalization, we performed chromatin immunoprecipitation (ChIP) assays using transgenic lines expressing a functional GFP-tagged *LHP1*. During vernalization, *LHP1* association with *FLC* chromatin increases and the association is maintained in warm conditions (Fig. 2), supporting a direct *in vivo* role for *LHP1* in the formation of repressive chromatin at *FLC*. Regions of *FLC* chromatin



**Figure 2** Role of *LHP1* in *FLC* repression. **(a)** Relative positions of amplified *FLC* regions used in the ChIP assay. **(b)** ChIP assay using antibody against GFP and chromatin from transgenic plants carrying GFP-tagged *LHP1* with and without vernalization. Three regions were examined for *LHP1* enrichment in *FLC* chromatin **(a)**. *AP3* was used for a loading control, because *AP3* is also repressed by *LHP1* (ref. 15) and its repression is not affected by vernalization **(c)**. **(c)** RT-PCR analysis of *AP3* expression in the *lhp1* mutant with and without vernalization.



**Figure 3** ChIP analyses of *FLC* chromatin. **(a)** H3K9 dimethylation levels of *FLC* chromatin before, during and after vernalization. **(b)** H3K4 trimethylation levels of *FLC* chromatin before, during and after vernalization. WT, wild-type.

enriched in LHP1 by vernalization (**Fig. 2b**) correspond to regions in which vernalization results in increased H3K9 dimethylation<sup>3</sup>.

In animals, HP1 interacts with the Suvar3-9 class of histone methyltransferases to maintain the methylated state of H3K9 in newly assembled nucleosomes during cell division<sup>12</sup>. In *lhp1* mutants, the vernalization-mediated dimethylation of *FLC* chromatin on H3K9 is not maintained (**Fig. 3**). Notably, H3K9 dimethylation at *FLC* still occurs in the absence of *LHP1* during a vernalizing cold exposure (**Fig. 3a**). Thus, *LHP1* is not required for the initiation of H3K9 dimethylation, but is essential for maintenance of H3K9 dimethylation and the silenced state of *FLC* after plants are returned to warm conditions.

In *D. melanogaster*, PcG and Trithorax group proteins act on target genes by association with specific DNA regulatory elements referred to as Polycomb and Trithorax response elements (PREs and TREs)<sup>8</sup>. A 2.8-kb region of the first intron of *FLC* is required for stable maintenance of *FLC* repression<sup>23</sup>. We find that a deletion of just 289 bp is sufficient to prevent maintenance of the silenced state of *FLC* (**Fig. 4a**; region labeled VRE for vernalization response element). Specifically, a deletion of this region in the first intron does not compromise *FLC* activation, nor does it prevent the repression of *FLC* expression during cold exposure (**Fig. 4b**); however, the VRE deletion prevents *FLC* repression from being maintained after a return to warm conditions (**Fig. 4b**) and it prevents vernalization from causing more rapid flowering (**Fig. 4c**).

Unlike the case in *lhp1* mutants in which H3K9 dimethylation increases only transiently during cold exposure, there is no increase in H3K9 dimethylation of *FLC* chromatin in the VRE deletion (**Fig. 3a**). Thus, the VRE region is required for the initial vernalization-mediated increase in H3K9 dimethylation. Whether or not the same H3K9 histone methyltransferase is involved in modifying *FLC* chromatin during and after cold exposure remains to be determined. Given that some of the proteins involved in the initiation versus maintenance phases of vernalization are different, and that there are 15 SET domain proteins that are predicted to methylate H3K9 in *A. thaliana*<sup>24</sup>, it is possible that different histone methyltransferases are involved at different stages of vernalization.


The intronic DNA component defined by the VRE deletion may act in vernalization-mediated *FLC* repression similar to a PRE in PcG-mediated repression. The binding of the *D. melanogaster* Dorsal

switch protein 1 (DSP1) to a minimal PRE can initiate recruitment of chromatin-remodeling complexes, including PRC2 and PRC1 (ref. 25). Similarly, the VRE region may serve as a site of recruitment of a vernalization-specific, chromatin-remodeling complex that is required for the initial increase in H3K9 dimethylation. Elevated H3K9 dimethylation alone is not sufficient to recruit LHP1, because not all chromatin with elevated levels of H3K9 dimethylation is associated with LHP1 (for instance, chromocenters; **Supplementary Figure 1**); therefore, additional factors, perhaps associated with the VRE, are required for LHP1 recruitment and stable silencing of euchromatic genes.

The failure to maintain *FLC* repression and the lack of a vernalization-mediated increase in H3K9 methylation at *FLC* chromatin in the VRE deletion are similar to the phenotypes of *vrn1* and *vrn2* mutants<sup>3,4</sup>. As in *vrn1* and *vrn2*, the initial repression of *FLC* during cold exposure occurs in the VRE deletion (**Fig. 4c**), indicating that this initial repression is mediated by other *cis* elements in *FLC* and does not involve increased H3K9 dimethylation.

The presence of *FRI*, which activates *FLC*, results in a higher level of H3K4 trimethylation at *FLC* chromatin than when *FRI* is absent<sup>26</sup>. During vernalization of *FRI*-Col, however, the level of H3K4 trimethylation in *FLC* chromatin is decreased, and this decrease is maintained after vernalization (**Fig. 3b**). Notably, in *lhp1* mutants the level of H3K4 trimethylation is also decreased during vernalization, but the reduced level is not maintained after plants are returned to warm conditions (**Fig. 3b**). Thus, *LHP1* is required to maintain the low level of H3K4 methylation associated with vernalization-mediated *FLC* silencing, but *LHP1* is not required for the initial decrease of H3K4 trimethylation during cold exposure. Similar to the *lhp1* mutant, trimethylated H3K4 levels at *FLC* chromatin are transiently reduced in the chromatin of the VRE deletion (**Fig. 3b**).

In animals, H3K4 trimethylation and active transcription of certain genes is mediated by a number of chromatin-remodeling proteins, including the PAF1 complex, which recruits the Trithorax group gene ASH1-like H3K4 methyltransferase<sup>27</sup>. PcG proteins act antagonistically to PAF1 and Trithorax group genes to maintain gene repression<sup>8</sup>. In *A. thaliana*, *FLC* regulation also seems to be mediated by antagonistic chromatin remodeling complexes. Active transcription and H3K4 methylation of *FLC* requires a PAF1 homolog<sup>26</sup> as well as an



**Figure 4** A region in the first intron of *FLC* is essential for maintenance of the vernalized state. **(a)** Location of an intronic region (VRE) required for maintenance. **(b)** RNA blot analyses of *FLC* expression before, during and after vernalization in lines described in **a** and **c**. **(c)** Flowering behavior of lines carrying either a full-length *FLC* construct or the construct with a VRE deletion in an *FRI/flc-3* background. Vernalization response of representative homozygous T3 lines.

ASH1-related H3K4 methyltransferase<sup>28</sup>. After a vernalizing cold exposure, the maintenance of the silenced state of *FLC* requires the association of LHP1 with *FLC* chromatin (Fig. 2b). This association might help to prevent complexes like PAF1 from activating *FLC* expression. Because *FLC* expression and H3K4 trimethylation transiently decrease during exposure to cold in *lhp1* mutants, the initial *FLC* repression system may involve other repressive complexes that compete with activating complexes.

In summary, vernalization-mediated epigenetic repression of *FLC* requires LHP1 as well as an intronic DNA component, and vernalization results in an increase in the localization of LHP1 at *FLC* chromatin. It is likely that LHP1 has a similar direct role in the PRC2-type repression of other genes in *A. thaliana* such as the floral homeotic gene *AGAMOUS* (*AG*)<sup>15</sup>, which is repressed by the *E(z)* homolog *CLF*<sup>9</sup>. A common theme for epigenetically stable euchromatic gene repression in both plants and other organisms is the 'histone code hypothesis'—a requirement for specific histone modifications and binding of proteins that recognize such modifications<sup>29</sup>. However, in the absence of PRC1 in plants, LHP1 may have a central role in the creation of mitotically stable repression of euchromatic genes.

## METHODS

**Plant materials.** We used the terminal flower2 mutants *tf12-1* and *tf12-2* (hereafter referred to as *lhp1-3* and *lhp1-4*, respectively)<sup>18</sup>. *lhp1-3* and *lhp1-4* in Columbia were introgressed into *FRI-Col*<sup>19</sup>.

**Vernalization treatment.** Seeds were germinated on agar plates for 5 d at 22 °C and vernalized for 40 d at 4 °C under 8 h of light and 16 h of dark. Post-vernalization samples continued to grow on agar plates under 8 h of light and 16 h of dark at 22 °C. All plants samples were prepared at 10 a.m. (2 h after light was turned on).

**ChIP analysis.** Constructs carrying LHP1::GFP were transformed into *lhp1-4* in *FRI-Col*, and homozygous lines were identified at T3 generation and used for ChIP analysis. Independently, *lhp1-4* in *Col* lines that carry LHP1::GFP were crossed to *FRI-Col* and F3 lines that are homozygous for both *FRI* and LHP1::GFP were used to confirm ChIP analysis. GFP antibodies were purchased from Molecular Probes. Antibodies against dimethylated H3K9 were purchased from Upstate. Both antibodies produced similar results. For biotin-conjugated antibodies, streptavidin agarose (Upstate) was used to precipitate immune complexes. ChIP assays were performed using a ChIP assay kit from Upstate following the manufacturer's suggested protocol, modified as previously described<sup>30</sup>. To detect the *FLC-P* region, we used a pair of primers that do not amplify a fragment from the *flc-3* deletion allele<sup>5</sup> (for primer sequences, see Supplementary Table 1 online). *FLC-V*, *FLC-U*, and *AP3* were amplified (for primer sequences, see Supplementary Table 1). The amount of chromatin was calculated as a multiple of the control using ImageQuant. All ChIP assays were performed at least three times from at least two biological replicates and produced similar results.

**RNA analysis.** Total RNA was isolated using TRI reagent (Sigma-Aldrich) according to the manufacturer's instructions. First-strand cDNA synthesis was performed on 2 µg of RNA using the M-MLV System for RT-PCR (Promega) followed by PCR amplification with ExTaq DNA Polymerase (Takara Mirus) according to the manufacturer's recommendations. *FLC*, *UBIQUITIN*, *FT* and *AP3* were amplified (for primer sequences, see Supplementary Table 1 online). Amplified fragments were separated on a 3% or 1.5% agarose gel. For RNA blot analysis, 15 µg of total RNA was separated on denaturing gel and transferred to a nylon membrane. The 3'-UTR of *FLC* was used as a probe. 18S RNA was visualized on the gel by ethidium bromide staining.

Note: Supplementary information is available on the Nature Genetics website.

## ACKNOWLEDGMENTS

We are grateful to S. Woody, R. Schmitz and R. Rodman for their comments on the manuscript. Research in the laboratory of R.M.A. was supported by the

College of Agricultural and Life Sciences of the University of Wisconsin and by grants from the US Department of Agriculture National Research Initiative Competitive Grants Program and the National Science Foundation. Research in the laboratory of S.E.J. was supported by US National Institute of Health grant GM060398.

## COMPETING INTERESTS STATEMENT

The authors declare that they have no competing financial interests.

Published online at <http://www.nature.com/naturegenetics>


Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>

1. Michaels, S.D. & Amasino, R.M. Memories of winter: vernalization and the competence to flower. *Plant Cell Environ.* **23**, 1145–1154 (2000).
2. Sung, S. & Amasino, R.M. REMEMBERING WINTER: Toward a molecular understanding of vernalization. *Annu. Rev. Plant Biol.* **56**, 491–508 (2005).
3. Sung, S. & Amasino, R.M. Vernalization in *Arabidopsis thaliana* is mediated by the PHD finger protein VIN3. *Nature* **427**, 159–164 (2004).
4. Bastow, R. *et al.* Vernalization requires epigenetic silencing of *FLC* by histone methylation. *Nature* **427**, 164–167 (2004).
5. Michaels, S.D. & Amasino, R.M. FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* **11**, 949–956 (1999).
6. Sheldon, C.C. *et al.* The *FLF* MADS box gene: a repressor of flowering in *Arabidopsis* regulated by vernalization and methylation. *Plant Cell* **11**, 445–458 (1999).
7. Gendall, A.R., Levy, Y.Y., Wilson, A. & Dean, C. The VERNALIZATION 2 gene mediates the epigenetic regulation of vernalization in *Arabidopsis*. *Cell* **107**, 525–535 (2001).
8. Ringrose, L. & Paro, R. Epigenetic regulation of cellular memory by the Polycomb and Trithorax group proteins. *Annu. Rev. Genet.* **38**, 413–443 (2004).
9. Schubert, D., Clarenz, O. & Goodrich, J. Epigenetic control of plant development by Polycomb-group proteins. *Curr. Opin. Plant Biol.* **8**, 553–561 (2005).
10. Gaudin, V. *et al.* Mutations in LIKE HETEROCHROMATIN PROTEIN 1 affect flowering time and plant architecture in *Arabidopsis*. *Development* **128**, 4847–4858 (2001).
11. Kotake, T., Takada, S., Nakahigashi, K., Ohto, M. & Goto, K. *Arabidopsis* TERMINAL FLOWER 2 gene encodes a heterochromatin protein 1 homolog and represses both FLOWERING LOCUS T to regulate flowering time and several floral homeotic genes. *Plant Cell Physiol.* **44**, 555–564 (2003).
12. Maison, C. & Almouzni, G. HP1 and the dynamics of heterochromatin maintenance. *Nat. Rev. Mol. Cell Biol.* **5**, 296–304 (2004).
13. Liu, L.P., Ni, J.Q., Shi, Y.D., Oakeley, E.J. & Sun, F.L. Sex-specific role of *Drosophila melanogaster* HP1 in regulating chromatin structure and gene transcription. *Nat. Genet.* **37**, 1361–1366 (2005).
14. Nielsen, S.J. *et al.* Rb targets histone H3 methylation and HP1 to promoters. *Nature* **412**, 561–565 (2001).
15. Nakahigashi, K., Jasencakova, Z., Schubert, I. & Goto, K. The *Arabidopsis* HETEROCHROMATIN PROTEIN1 homolog (TERMINAL FLOWER2) silences genes within euchromatic region but not genes positioned in heterochromatin. *Plant Cell Physiol.* **46**, 1747–1756 (2005).
16. Libault, M. *et al.* The *Arabidopsis* LHP1 protein is a component of euchromatin. *Planta* **222**, 910–925 (2005).
17. Zemach, A. *et al.* Different domains control the localization and mobility of LIKE HETEROCHROMATIN PROTEIN1 in *Arabidopsis* nuclei. *Plant Cell* **18**, 133–145 (2006).
18. Lindroth, A.M. *et al.* Dual histone H3 methylation marks at lysines 9 and 27 required for interaction with CHROMOMETHYLASE3. *EMBO J.* **23**, 4286–4296 (2004).
19. Lee, I. & Amasino, R.M. Effect of vernalization, photoperiod, and light quality on the flowering phenotype of *Arabidopsis* plants containing the FRIGIDA gene. *Plant Physiol.* **108**, 157–162 (1995).
20. Mylne, J.S. *et al.* LHP1, the *Arabidopsis* homologue of HETEROCHROMATIN PROTEIN1, is required for epigenetic silencing of *FLC*. *Proc. Natl. Acad. Sci. USA* **103**, 5012–5017 (2006).
21. Valverde, F. *et al.* Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. *Science* **303**, 1003–1006 (2004).
22. Huang, T., Bohlenius, H., Eriksson, S., Parcy, F. & Nilsson, O. The mRNA of the *Arabidopsis* gene *FT* moves from leaf to shoot apex and induces flowering. *Science* **309**, 1694–1696 (2005).
23. Sheldon, C.C., Conn, A.B., Dennis, E.S. & Peacock, W.J. Different regulatory regions are required for the vernalization-induced repression of FLOWERING LOCUS C and for the epigenetic maintenance of repression. *Plant Cell* **14**, 2527–2537 (2002).
24. Baumbusch, L.O. *et al.* The *Arabidopsis thaliana* genome contains at least 29 active genes encoding SET domain proteins that can be assigned to four evolutionarily conserved classes. *Nucleic Acids Res.* **29**, 4319–4333 (2001).
25. Dejardin, J. *et al.* Recruitment of *Drosophila* Polycomb group proteins to chromatin by DSP1. *Nature* **434**, 533–538 (2005).

26. He, Y., Doyle, M.R. & Amasino, R.M. PAF1-complex-mediated histone methylation of FLOWERING LOCUS C chromatin is required for the vernalization-responsive, winter-annual habit in *Arabidopsis*. *Genes Dev.* **18**, 2774–2784 (2004).
27. Krogan, N.J. *et al.* The Paf1 complex is required for histone H3 methylation by COMPASS and Dot1p: linking transcriptional elongation to histone methylation. *Mol. Cell* **11**, 721–729 (2003).
28. Kim, S.Y. *et al.* Establishment of the vernalization-responsive, winter-annual habit in *Arabidopsis* requires a putative histone H3 methyl transferase. *Plant Cell* **17**, 3301–3310 (2005).
29. Jenuwein, T. & Allis, C.D. Translating the histone code. *Science* **293**, 1074–1080 (2001).
30. Johnson, L., Cao, X. & Jacobsen, S. Interplay between two epigenetic marks. DNA methylation and histone H3 lysine 9 methylation. *Curr. Biol.* **12**, 1360–1367 (2002).

**Supplementary Figure 1.** LHP1 localization.

GFP-tagged LHP1 (upper panel) and FLAG-tagged LHP1 (lower panel) are localized in euchromatic regions. Chromocenters are visualized by either DAPI staining and immuno-staining using antibody against dimethylated histone H3K9.



*lhp1-4* in Col plants were transformed with LHP1::GFP or LHP1::FLAG and homozygous T3 lines were used for analyses. Both the GFP-tagged and FLAG-tagged LHP1 constructs can rescue the *lhp1* mutant phenotype. Interphase nuclei were isolated from leaves of 3 week old plants and immunolabeling was performed as previously described<sup>1</sup>. Histone H3 2MK9 was detected using polyclonal antisera from Upstate Biotechnology (#07-441, diluted 1/150) and FLAG was detected using monoclonal antibodies from Sigma (#F 3165, diluted 1/100). Rhodamine Red

conjugated antibodies from Jackson Immunoresearch (#111 295 144, goat anti-rabbit diluted 1/200) and FITC conjugated antibodies from Abcam (#ab7064, goat anti-mouse diluted 1/200) were used for secondary antibodies. GFP was observed directly. After staining, vectashield mounting medium with DAPI was added and nuclei were visualized using a Zeiss Axioimager Z1 motorized microscope with Apotome system equipped with ORCA-ER camera.

1. Jasencakova, Z. et al. Histone modifications in Arabidopsis- high methylation of H3 lysine 9 is dispensable for constitutive heterochromatin. *Plant J* **33**, 471-80 (2003).

**Supplementary Table 1. Primer sequences used for ChIP and RT-PCR analyses.**

<b>Primers used for ChIP analyses</b>	
FLC-P	5'-cgacttgaacccaacacctgaggatcaaat-3'
	5'-agaagataaaaggggaacaatgaaaag-3'
FLC-V	5'- ttattatagatccgtaccaagagggtg-3'
	5'- atagattgcctcatattatgtgattg-3'
FLC-U	5'- ttgcaaaataagccgtaggcttcttctact-3'
	5'-cgctgataaggcgagcgtttgtatatct-3'
AP3	5'- gaagagattggtggagaggacaagagat-3'
	5'-atcacttagtttcatcaacttctgaact-3'
<b>Primers used for RT-PCR analyses</b>	
FLC	5'-ttctcaaactcgcaacggtctc-3'
	5'-gattgtccagcaggtgacatctc-3'
FT	5'- tacgaaaatccaagtcccactg-3'
	5'- aaactcgcgagtgttgaagttc-3'
AP3	5'- aagtagttactaaataattagatagacaa-3'
	5'- tcgaagggtcacgtgcttacgctctctg-3'
UBQ	5'-gatcttgcggaaaacaattggaggatgg-3'
	5'-cgacttgcattagaagaagagataacagg-3'